

11/9/11 – Linear Algebra and Abstract Algebra

1. Let d_n be the determinant of the $n \times n$ matrix whose entries, from left to right and then from top to bottom, are $\cos 1, \cos 2, \dots, \cos n^2$. (For example,

$$d_3 = \begin{vmatrix} \cos 1 & \cos 2 & \cos 3 \\ \cos 4 & \cos 5 & \cos 6 \\ \cos 7 & \cos 8 & \cos 9 \end{vmatrix}.$$

The argument of \cos is always in radians, not degrees.) Evaluate $\lim_{n \rightarrow \infty} d_n$. (Putnam 2009)

2. Alan and Barbara play a game in which they take turns filling entries of an initially empty 2008×2008 array. Alan plays first. At each turn, a player chooses a real number and places it in a vacant entry. The game ends when all the entries are filled. Alan wins if the determinant of the resulting matrix is nonzero; Barbara wins if it is zero. Which player has a winning strategy? (Putnam 2008)

3. Let H be an $n \times n$ matrix all of whose entries are ± 1 and whose rows are mutually orthogonal. Suppose H has an $a \times b$ submatrix whose entries are all 1. Show that $ab \leq n$. (Putnam 2005)

4. Let \mathbf{A} and \mathbf{B} be different $n \times n$ matrices with real entries. If $\mathbf{A}^3 = \mathbf{B}^3$ and $\mathbf{A}^2\mathbf{B} = \mathbf{B}^2\mathbf{A}$, can $\mathbf{A}^2 + \mathbf{B}^2$ be invertible? (Putnam 1991)

5. Let S be a non-empty set with an associative operation that is left and right cancellative ($xy = xz$ implies $y = z$, and $yx = zx$ implies $y = z$). Assume that for every a in S the set $\{a^n : n = 1, 2, 3, \dots\}$ is finite. Must S be a group? (Putnam 1989)

6. Let S be a set of real numbers which is closed under multiplication (that is, if a and b are in S , then so is ab). Let T and U be disjoint subsets of S whose union is S . Given that the product of any *three* (not necessarily distinct) elements of T is in T and that the product of any three elements of U is in U , show that at least one of the two subsets T, U is closed under multiplication. (Putnam 1995)

7. Let G be a group with identity e and $\phi : G \rightarrow G$ a function such that

$$\phi(g_1)\phi(g_2)\phi(g_3) = \phi(h_1)\phi(h_2)\phi(h_3)$$

whenever $g_1g_2g_3 = e = h_1h_2h_3$. Prove that there exists an element $a \in G$ such that $\psi(x) = a\phi(x)$ is a homomorphism (i.e. $\psi(xy) = \psi(x)\psi(y)$ for all $x, y \in G$). (Putnam 1997)