

Problem Set 8

March 28 and 29, in class

Recall some facts:

- If $f(x) \leq g(x)$ for all x in $[a, b]$ then $\int_a^b f(x) dx \leq \int_a^b g(x) dx$.
- If k is a constant then $\int_a^b kf(x) dx = k \int_a^b f(x) dx$.
- $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$.

1. (a) Understand the following proof:

$$\begin{aligned} -|f(x)| &\leq f(x) \leq |f(x)| \\ -\int_a^b |f(x)| dx &\leq \int_a^b f(x) dx \leq \int_a^b |f(x)| dx \\ \left| \int_a^b f(x) dx \right| &\leq \int_a^b |f(x)| dx \end{aligned}$$

- (b) Explain why the last inequality makes sense. Hint: Find some functions for which the left-hand side and the right-hand side are not equal. Then explain why the left-hand side always has to be less.

2. Suppose that f is continuous, $a < b$, and $\int_a^b f(x) dx = 0$.

- (a) Does it necessarily follow that $f(x) = 0$ for all x in $[a, b]$?
- (b) Does it necessarily follow that $f(x) = 0$ for *some* x in $[a, b]$?
- (c) Does it necessarily follow that $\int_a^b |f(x)| dx = 0$?
- (d) Does it necessarily follow that $\left| \int_a^b f(x) dx \right| = 0$?
- (e) Must all the upper sums $U(P)$ be nonnegative?
- (f) Must all the upper sums $U(P)$ be positive?
- (g) Can a lower sum $L(P)$ be positive?

3. Suppose that f and g are continuous, $a < b$, and $\int_a^b f(x) dx \geq \int_a^b g(x) dx$.

(a) Does it necessarily follow that $\int_a^b [f(x) - g(x)] dx \geq 0$?

(b) Does it necessarily follow that $f(x) \geq g(x)$ for all x in $[a, b]$?

(c) Does it necessarily follow that $f(x) \geq g(x)$ for *some* x in $[a, b]$?

(d) Does it necessarily follow that $\left| \int_a^b f(x) dx \right| \geq \left| \int_a^b g(x) dx \right|$?

(e) Does it necessarily follow that $\int_a^b |f(x)| dx \geq \int_a^b |g(x)| dx$?

(f) Does it necessarily follow that $\int_a^b |f(x)| dx \geq \int_a^b g(x) dx$?

4. While $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$, in general it is **not true** that

$$\int_a^b f(x)g(x) dx = \left(\int_a^b f(x) dx \right) \left(\int_a^b g(x) dx \right).$$

(a) Find two numbers a and b and two functions f and g for which the equation above fails—any random choice of a , b , f , and g should work.

(b) Find two numbers a and b and two functions f and g for which the equation above holds—you will have to choose a , b , f , and g carefully.