

The Pfaffian-Grassmannian derived equivalence

Lev Borisov, Andrei Căldăraru*

Abstract

We argue that there exists a derived equivalence between Calabi-Yau threefolds obtained by taking dual hyperplane sections (of the appropriate codimension) of the Grassmannian $\mathbf{G}(2, 7)$ and the Pfaffian $\mathbf{Pf}(7)$. The existence of such an equivalence has been conjectured by physicists for almost ten years, as the two families of Calabi-Yau threefolds are believed to have the same mirror. It is the first example of a derived equivalence between Calabi-Yau threefolds which are provably non-birational.

Introduction

0.1. Let V be a vector space of dimension seven over \mathbf{C} (or any algebraically closed field of characteristic zero), and let

$$\mathbf{G} = \mathbf{G}(2, V)$$

be the Grassmannian of planes in V . The Plücker map embeds \mathbf{G} as a smooth subvariety of dimension 10 of

$$\mathbf{P} = \mathbf{P}^{20} = \mathbf{P}(\wedge^2 V).$$

Regard the dual projective space

$$\mathbf{P}^* = \mathbf{P}(\wedge^2 V^*)$$

as the projectivization of the space of two-forms on V . The Pfaffian locus

$$\mathbf{Pf} \subset \mathbf{P}^*$$

is defined to be the projectivization of the locus of degenerate two-forms on V . Equations for \mathbf{Pf} can be obtained by taking the Pfaffians of the diagonal minors of a skew-symmetric 7×7 matrix of linear forms on V .

While the Grassmannian \mathbf{G} is smooth, the Pfaffian \mathbf{Pf} is a singular subvariety of \mathbf{P}^* of dimension 17. Indeed, a point $\omega \in \mathbf{Pf}$ will be singular precisely when the rank of ω is two. (Recall that the rank of a two-form ω – or, equivalently, of a skew symmetric matrix – is always even. In our case, since we eliminate $\omega = 0$ by projectivizing, this rank could be two, four, or six. The general two-form on V has rank six; it has rank four at the smooth points of \mathbf{Pf} , and it has rank two at its singular points, which coincide with $\mathbf{G}(2, V^*)$ in its Plücker embedding.)

The Pfaffian is the classical projective dual of the Grassmannian:

$$\mathbf{Pf} = \{\mathbf{y} \in \mathbf{P}^* : \mathbf{G} \cap H_{\mathbf{y}} \text{ is singular}\},$$

where $H_{\mathbf{y}}$ is the hyperplane in \mathbf{P} corresponding to \mathbf{y} . See, for example, Proposition 1.5.

*Mathematics Department, University of Wisconsin–Madison, 480 Lincoln Drive, Madison, WI 53706–1388, USA, *e-mail*: andreic@math.wisc.edu, borisov@math.wisc.edu

0.2. Consider a seven-dimensional linear subspace

$$W \subset \wedge^2 V^*,$$

and denote by \mathbf{W} its image in \mathbf{P}^* . Let Y be the intersection of \mathbf{W} with \mathbf{Pf} .

On the dual side, let

$$\mathbf{M} = \text{Ann}(W) \subset \wedge^2 V$$

be the 14-dimensional annihilator of W ; again, we will use bold-face \mathbf{M} to denote its projectivization in \mathbf{P} , which has codimension seven. Let X be the intersection of \mathbf{M} and \mathbf{G} .

The main result of this paper is the following theorem.

0.3. Theorem. *For a given choice of W , if either X or Y has dimension three, then X is smooth if and only if Y is. When this happens, X and Y are Calabi-Yau threefolds with*

$$h^{1,1} = 1, \quad h^{1,2} = 50,$$

and there exists an equivalence of derived categories

$$\Phi : \mathbf{D}_{\text{coh}}^b(Y) \xrightarrow{\sim} \mathbf{D}_{\text{coh}}^b(X).$$

0.4. Such a result has been conjectured for a while. Indeed, Rødland [9] argued, by comparing solutions to the Picard-Fuchs equation, that the families of X 's and of Y 's appear to have the same mirror family. Recently, Hori and Tong [4] gave a more detailed string theory argument supporting the same conclusion. If we denote by Z their common mirror, Kontsevich's Homological Mirror Symmetry conjecture predicts

$$\mathbf{D}_{\text{coh}}^b(X) \cong \mathbf{Fuk}(Z) \cong \mathbf{D}_{\text{coh}}^b(Y).$$

0.5. This is the first example of a derived equivalence between Calabi-Yau threefolds which can be proved to be non-birational. Indeed, if X and Y were birational, they would have to differ by a sequence of projective flops because they are minimal in the sense of Mori theory ([5, 6.38]). On the other hand, no such flops are possible on either X or Y , because their Picard numbers are $h^{1,1} = 1$, and hence all their non-zero nef divisors are ample. Therefore they would have to be isomorphic. However, this can not be true, since if we denote by H_X and H_Y , respectively, the ample generators of the Picard groups of X and of Y , we have [9]

$$H_X^3 = 42, \quad H_Y^3 = 14.$$

0.6. Our results appear to fit very well with the theory of Homological Projective Duality developed by Kuznetsov [6]. Indeed, we have a pair of varieties \mathbf{G} and \mathbf{Pf} , embedded in dual projective spaces, whose dual hyperplane sections are derived equivalent. It would be interesting to understand this relationship further. To this end, we make some comments in Section 7 about how this example seems to fit in the general theory. It is worth noting that Kuznetsov has independently arrived at a proof of Theorem 0.3, see [8].

0.7. Let us now briefly describe the construction of the derived equivalence Φ . Recall [1] that, in order to give Φ , it is essentially enough to describe the image of Φ on structure sheaves of points \mathcal{O}_y for $y \in Y$, and to check that the family $\{\Phi\mathcal{O}_y\}_{y \in Y}$ is an orthonormal basis for $\mathbf{D}_{\text{coh}}^b(X)$.

A point y in Y can be regarded as a two-form on V of rank four. As such it has a kernel K which is a three-dimensional linear subspace of V . On the other hand, a point x in X corresponds to a two-dimensional linear subspace T of V . For a general choice of x and y we will have

$$T \cap K = 0.$$

However, for a fixed $y \in Y$, the set of points $x \in X$ for which T intersects K non-trivially is a curve C_y in X . Our choice for the functor Φ is to set

$$\Phi\mathcal{O}_y = \mathcal{I}_{C_y},$$

where \mathcal{I}_{C_y} denotes the ideal sheaf of the curve C_y . The technical core of the paper consists then of showing that this choice gives rise to an orthonormal family.

It would be interesting to find out the relationship of this approach to the construction of Donaldson-Thomas moduli spaces. In the course of our proof we see that Y is such a moduli space of ideal sheaves of curves on X . Perhaps other examples of derived equivalences can be obtained as Donaldson-Thomas spaces.

0.8. The paper is organized as follows. In Section 1 we set up the notation, and we recall the appropriate definitions from linear algebra. We also sketch the standard constructions of the tangent spaces to the Pfaffian and Grassmannian varieties. In Section 2 we give a purely linear-algebra argument for the fact that X is smooth if and only if Y is. We study the geometry of certain Schubert cycles in Section 3, and we prove a vanishing result for Ext groups between them on the Grassmannian. In Section 4 we define the curves C_y which are parametrized by the points of Y , and we argue that they all have dimension one. Section 5 reduces the computation of the orthogonality of the family $\{\mathcal{I}_{C_y}\}$ to the vanishing result on \mathbf{G} proved earlier. We define the functor Φ in Section 6 and we argue that it is an equivalence. Finally, in Section 7 we discuss connections with results of Kuznetsov on Homological Projective Duality.

0.9. Acknowledgments. The second author would like to thank Mark Gross for first introducing him to the physicists' conjecture, to Kentaro Hori for pointing out the reference [9], and to Ron Donagi for comments on the geometry of the Grassmannian. Michael Stillman provided comments on the initial Macaulay computations which confirmed our results. Dan Abramovich had comments on the relationship to Donaldson-Thomas invariants. Alexander Kuznetsov explained connections with Homological Projective Duality in higher dimensions, and pointed out an alternative argument for \mathbf{G} -vanishing. This material is based upon work supported by the National Science Foundation under Grants No. DMS-0456801 and DMS-0556042.

1. Basic facts from linear algebra

In this section we set up the basic notation and review some standard results from linear algebra and differential geometry.

1.1. Let V be a vector space. Denote by $\mathbf{G}(k, V)$ the Grassmannian of k -dimensional planes in V . The Plücker embedding is the map

$$\mathbf{G}(k, V) \hookrightarrow \mathbf{P}(\wedge^k V),$$

which maps a point x of $\mathbf{G}(k, V)$ (i.e., a k -plane $x \subset V$) to the point

$$[e_1 \wedge e_2 \wedge \cdots \wedge e_k] \in \mathbf{P}(\wedge^k V),$$

where e_1, \dots, e_k is any basis of x .

If x is a point in the Grassmannian, the tangent space $T_{\mathbf{G}(k, V), x}$ is canonically identified with

$$\text{Hom}(x, V/x).$$

Thus, if V has dimension n , the Grassmannian $\mathbf{G}(k, V)$ has dimension $k(n - k)$.

In particular, if $k = 1$, the tangent space $T_{\mathbf{P}V, x}$ to the projective space at a point x is naturally identified with $\text{Hom}(x, V/x)$ which is (up to scalars) the same as V/x .

1.2. The tangent map to the Plücker embedding is the map

$$\text{Hom}(x, V/x) \rightarrow \text{Hom}(\wedge^k x, \wedge^k V / \wedge^k x),$$

given by

$$\varphi \mapsto \left(v_1 \wedge \cdots \wedge v_k \mapsto \sum_{i=1}^k v_1 \wedge v_2 \wedge \cdots \wedge v_{i-1} \wedge \varphi(v_i) \wedge v_{i+1} \wedge \cdots \wedge v_k \right).$$

1.3. An element $y \in \wedge^k V^*$ will be called a k -form on V . Since we have

$$\wedge^k V^* \cong (\wedge^k V)^*,$$

the annihilator of y , $\text{Ann}(y)$, is a hyperplane $H \subset \wedge^k V$. It consists of all $x \in \wedge^k V$ such that $y(x) = 0$.

1.4. From now on we will concentrate on the case $k = 2$, and define

$$\mathbf{P} = \mathbf{P}(\wedge^2 V), \quad \mathbf{P}^* = \mathbf{P}(\wedge^2 V^*), \quad \mathbf{G} = \mathbf{G}(2, V).$$

Frequently, when regarding a point $x \in \mathbf{G}$ as a two-plane in V we will call this plane T .

If y is a two-form on V , its kernel K is defined to be the set of all $v \in V$ such that $y(v \wedge w) = 0$ for all $w \in V$. We define the rank of y by

$$\text{rk } y = n - \dim K.$$

The rank of a two-form is always even.

The kernel of a form y does not change if we multiply y by a non-zero scalar. Thus we can speak of the kernel of a point $y \in \mathbf{P}^*$, and we will frequently refer to such a point as a two-form.

1.5. Proposition. *Let \mathbf{y} be a point in \mathbf{P}^* , and let H be the corresponding hyperplane in \mathbf{P} . Let x be a point of intersection of H and \mathbf{G} . Then H is tangent to \mathbf{G} at x (under the Plücker embedding) if and only if $T \subset K$, where T is the two-plane in V corresponding to x , and K is the kernel of \mathbf{y} , regarded as a two-form on V .*

Proof. First assume $T \subset K$. Let φ be a tangent vector to \mathbf{G} at x , i.e., a morphism $\varphi : T \rightarrow V/T$. Then the image of φ under the differential of the Plücker map is the map $\wedge^2 T \rightarrow \wedge^2 V / \wedge^2 T$ given by

$$\mathbf{t}_1 \wedge \mathbf{t}_2 \mapsto \varphi(\mathbf{t}_1) \wedge \mathbf{t}_2 + \mathbf{t}_1 \wedge \varphi(\mathbf{t}_2).$$

Choosing \mathbf{t}_1 and \mathbf{t}_2 linearly independent vectors in T , we identify this (up to scalars) with the element

$$\mathbf{t} = \varphi(\mathbf{t}_1) \wedge \mathbf{t}_2 + \mathbf{t}_1 \wedge \varphi(\mathbf{t}_2)$$

of $\wedge^2 V / \wedge^2 T$. Since $T \subset K$, $\mathbf{t}_1, \mathbf{t}_2$ are in the kernel of \mathbf{y} , and therefore \mathbf{y} vanishes on \mathbf{t} . In other words, we have proven that

$$\varphi \in T_{\mathbf{G},x} \implies \varphi \in T_{H,x},$$

or that H is tangent to \mathbf{G} at x .

Conversely, assume $T \not\subset K$. At least one of $\mathbf{t}_1, \mathbf{t}_2$ is not in K , assume it is \mathbf{t}_1 . There exists then $\mathbf{v} \in V$ such that $\mathbf{y}(\mathbf{t}_1 \wedge \mathbf{v}) \neq 0$. Define φ on the basis $\mathbf{t}_1, \mathbf{t}_2$ of T by setting

$$\begin{aligned} \varphi(\mathbf{t}_1) &= 0 \\ \varphi(\mathbf{t}_2) &= \mathbf{v}. \end{aligned}$$

Then $\mathbf{y}(\mathbf{t}) \neq 0$, and therefore the tangent vector to \mathbf{G} at x corresponding to φ is not in H . We conclude that H is not tangent to \mathbf{G} at x . \square

1.6. From now on assume that $n = \dim V$ is odd. The Pfaffian locus $\mathbf{Pf} \subset \mathbf{P}^*$ is defined to be the locus of non-zero degenerate two-forms on V . (A form is called degenerate if its rank is less than $n - 1$.) If we let A be a generic $n \times n$ matrix of linear forms on V , then \mathbf{Pf} is cut out by the n Pfaffians of $(n - 1) \times (n - 1)$ diagonal minors of A (obtained by removing the i -th row and column of A , for $i = 1, \dots, n$).

Since a point $\mathbf{y} \in \mathbf{Pf}$ corresponds to a two-form on V , its kernel is a subspace $K \subset V$. If V is odd-dimensional, \mathbf{y} being degenerate implies K is at least three-dimensional. Since we've eliminated the zero form by projectivizing, if $\dim V = 7$, $\dim K$ could only be three or five (the rank of \mathbf{y} can be either four or two).

1.7. Proposition 1.5 shows that the Pfaffian is precisely the classical dual variety to \mathbf{G} : indeed, $H \cap \mathbf{G}$ is singular for a hyperplane H if and only if H is tangent at some point of \mathbf{G} , i.e., as a point of \mathbf{P}^* , it corresponds to a form with kernel of dimension ≥ 3 .

1.8. Unlike the Grassmannian, which is smooth, the Pfaffian is singular at the points where the rank drops further (e.g., for $\dim V = 7$, when the rank is two). At singular points $\mathbf{y} \in \mathbf{Pf}$, we have $T_{\mathbf{Pf},\mathbf{y}} = T_{\mathbf{P}^*,\mathbf{y}}$. If \mathbf{y} is smooth, the following proposition describes the tangent space $T_{\mathbf{Pf},\mathbf{y}} \subset T_{\mathbf{P}^*,\mathbf{y}}$.

1.9. Proposition. *Let $\mathbf{y} \in \mathbf{Pf}$ correspond to a degenerate two-form on V with kernel $K \subset V$ of dimension three. Identify, up to scalars, $\mathbb{T}_{\mathbf{P}^*, \mathbf{y}}$ with $\wedge^2 V^* / \langle \mathbf{y} \rangle$, so that vectors in $\mathbb{T}_{\mathbf{P}^*, \mathbf{y}}$ are thought of as two-forms modulo \mathbf{y} . Then $v \in \mathbb{T}_{\mathbf{P}^*, \mathbf{y}}$ is tangent to \mathbf{Pf} at \mathbf{y} if and only if K is isotropic for v , i.e.,*

$$v(k_1 \wedge k_2) = 0 \text{ for all } k_1, k_2 \in K.$$

Proof. Pick a nonzero element $\mathbf{y}_0 \in \wedge^2 V^*$ in the cone over the Pfaffian that maps to \mathbf{y} under projectivization. Since all rank $n-3$ skew forms on V are in the same orbit of the $\mathrm{GL}(V)$ action, \mathbf{y}_0 is contained in the dense open subset of the cone over the Pfaffian which is its orbit under the $\mathrm{GL}(V)$ action. As a result, the tangent space to $\mathrm{Cone}(\mathbf{Pf})$ at \mathbf{y}_0 can be viewed as the span of $g\mathbf{y}_0$ for $g \in \mathrm{Lie}(\mathrm{GL}(V)) = \mathrm{End}(V^*)$ under the natural action. In an appropriate basis of V^* one has $\mathbf{y}_0 = x_1 \wedge x_2 + \dots + x_{n-4} \wedge x_{n-3}$, and $g\mathbf{y}_0 = gx_1 \wedge x_2 + x_1 \wedge gx_2 + \dots + gx_{n-4} \wedge x_{n-3} + x_{n-4} \wedge gx_{n-3}$. Since $x_i \in \mathrm{Ann}(K)$, we see, after passing from $\mathrm{Cone}(\mathbf{Pf})$ to \mathbf{Pf} , that the tangent space to \mathbf{Pf} is contained in the space of forms that make K isotropic. It is easy to see that the Pfaffian and this space are of the same dimension, which finishes the proof. \square

2. Simultaneous smoothness

From now on, the space V is assumed to be seven-dimensional. In this section we argue that, for a given choice of seven-dimensional linear subspace

$$W \subset \wedge^2 V^*,$$

the hyperplane sections

$$X = \mathbf{M} \cap \mathbf{G}$$

and

$$Y = \mathbf{W} \cap \mathbf{Pf}$$

are either both smooth, or both singular, at least in the case when X and Y are of the (expected) dimension three.

2.1. In the previous section we noted that a point $x \in X$ corresponds to a plane $T \subset V$, while a point $\mathbf{y} \in Y$, regarded as a two-form on V , has a kernel $K \subset V$, of dimension three or five.

Let $R \subset X \times Y$ denote the locus of pairs (x, \mathbf{y}) for which $T \subset K$. Let π_X, π_Y be the projections from R to X and to Y , respectively.

2.2. Proposition. *The following statements hold:*

- (a) *the set of points $x \in X$ where $\dim \mathbb{T}_{X, x} > 3$ coincides with the image $\pi_X(R)$;*
- (b) *the set of points $\mathbf{y} \in Y$ where $\dim \mathbb{T}_{Y, \mathbf{y}} > 3$ coincides with the image $\pi_Y(R)$.*

Proof. (a) Let $x \in \mathbf{G}$ be a point on the Grassmannian, and \mathbf{y} an arbitrary point in \mathbf{P}^* . Let H be the hyperplane in \mathbf{P} corresponding to \mathbf{y} . By Proposition 1.5, the following two statements are equivalent:

- H is tangent to \mathbf{G} at x ;

– $T \subset K = \text{Ker}(\mathbf{y})$.

Assume first that x is a point in X such that $\dim T_{X,x} > 3$. Then $T_{X,x}$, which is the intersection of $T_{\mathbf{G},x}$ and \mathbf{M} inside $T_{\mathbf{P},x}$, has dimension higher than the expected

$$3 = 10 + 13 - 20.$$

Therefore $T_{\mathbf{G},x}$ and \mathbf{M} do not span all of $T_{\mathbf{P},x}$, and there exists a hyperplane H inside $T_{\mathbf{P},x}$ (and thus a hyperplane in \mathbf{P} through x) containing both. Let \mathbf{y} be the point in \mathbf{P}^* corresponding to H .

Since H is tangent to \mathbf{G} at x , $T \subset K$ by the claim. Therefore $\mathbf{y} \in \mathbf{Pf}$ (because it has a kernel of dimension ≥ 2). Since we also have $\mathbf{M} \subset H$, it follows that $\mathbf{y} \in \mathbf{W}$, and therefore $\mathbf{y} \in Y$. The pair (x, \mathbf{y}) is thus in R , and x is in the image $\pi_X(R)$.

Conversely, assume that x in X is in the image of π_X , and let (x, \mathbf{y}) be a point in R . Let H be the hyperplane in \mathbf{P} corresponding to \mathbf{y} . It then follows that H contains \mathbf{M} (because $\mathbf{y} \in \mathbf{W}$), and therefore $x \in H$. The claim implies now that H is also tangent to \mathbf{G} at x , and therefore the tangent space to H at x contains $T_{\mathbf{G},x}$ and $T_{\mathbf{M},x}$. Thus the intersection of these two spaces (which is $T_{X,x}$) can not be of the expected dimension, and therefore $\dim T_{X,x} > 3$.

(b) Let \mathbf{y} be a point on Y . If the rank of \mathbf{y} , regarded as a two-form on V , is two, then $\dim T_{Y,\mathbf{y}} > 3$ because the Pfaffian is already singular at \mathbf{y} , and cutting it down by a codimension 14 hyperplane \mathbf{W} will not cut down the dimension of the tangent space to three. Thus we need to show that \mathbf{y} is in the image of R , in other words, that there exists a two dimensional space $T \subset K$ such that $x \in X$. The kernel K is five-dimensional, and thus $\mathbf{G}(2, K)$ is a subvariety of \mathbf{G} of dimension six, completely contained in the hyperplane in \mathbf{P} corresponding to \mathbf{y} . Completing $\{\mathbf{y}\}$ to a basis of W gives rise to six more hyperplanes in \mathbf{P} , which must have a common point x in $\mathbf{G}(2, K)$. Thus x is in X (being on \mathbf{G} and at the intersection of the seven hyperplanes corresponding to W), and $T \subset K$. It follows that $(x, \mathbf{y}) \in R$, thus \mathbf{y} is in the image of π_Y .

We can assume thus that \mathbf{y} is a smooth point of \mathbf{Pf} , the rank of \mathbf{y} is four, and thus its kernel $K \subset V$ has dimension three. The tangent space $T_{\mathbf{Pf},\mathbf{y}}$ consists of all the tangent vectors in $T_{\mathbf{P}^*,\mathbf{y}}$ for which K is isotropic (Proposition 1.9). Explicitly, we have

$$T_{\mathbf{P}^*,\mathbf{y}} \cong (\wedge^2 V^*) / \langle \mathbf{y} \rangle,$$

and a two-form $\omega \in \wedge^2 V^*$ will be tangent to \mathbf{Pf} at \mathbf{y} if and only if it vanishes on $\wedge^2 K$. In other words, it is (modulo \mathbf{y}) the subspace in $\wedge^2 V^*$ whose annihilator is $\wedge^2 K$.

The statement $\dim T_{Y,\mathbf{y}} > 3$ is equivalent to $T_{\mathbf{W},\mathbf{y}}$ and $T_{\mathbf{Pf},\mathbf{y}}$ not intersecting transversely in $T_{\mathbf{P}^*,\mathbf{y}}$, i.e., not spanning the full $T_{\mathbf{P}^*,\mathbf{y}}$. Since we have

$$\text{Ann}(T_{\mathbf{W},\mathbf{y}}) \cap \text{Ann}(T_{\mathbf{Pf},\mathbf{y}}) = \text{Ann}(\langle T_{\mathbf{W},\mathbf{y}}, T_{\mathbf{Pf},\mathbf{y}} \rangle),$$

it follows that this is equivalent to the existence of a non-zero

$$x \in \text{Ann}(T_{\mathbf{W},\mathbf{y}}) \cap \text{Ann}(T_{\mathbf{Pf},\mathbf{y}}) \cap \text{Ann}(\mathbf{y}).$$

Being in $\text{Ann}(T_{\mathbf{W},\mathbf{y}}) \cap \text{Ann}(\mathbf{y})$ is equivalent to being in \mathbf{M} , while being in $\text{Ann}(T_{\mathbf{Pf},\mathbf{y}})$ is equivalent to being a point on \mathbf{G} for which $T \subset K$. Thus $\dim T_{Y,\mathbf{y}} > 3$ is equivalent to the existence of an $x \in X$ such that $(x, \mathbf{y}) \in R$. \square

2.3. Corollary. *Assume either X or Y has dimension three. Then X is smooth if and only if Y is smooth.*

Proof. Both statements are equivalent to $R = \emptyset$. □

2.4. Remark. In [9], Rødland argues that $h^{1,1}(X) = h^{1,1}(Y) = 1$ and $h^{1,2}(X) = h^{1,2}(Y) = 50$ for generic cuts. In fact, this statement holds whenever X and Y are smooth: indeed, the family of such cuts is smooth over the appropriate open subset of $\mathbf{G}(7, \wedge^2 V^*)$, and thus all fibers are diffeomorphic.

3. A vanishing result for Schubert cycles

In this section we define a family $\{S_{\mathbf{y}}\}$ of Schubert cycles in \mathbf{G} , parametrized by the smooth points \mathbf{y} of \mathbf{Pf} , and we study homological orthogonality properties of this family. The most important result is \mathbf{G} -vanishing, which is the statement that for $0 \leq j \leq 5$, and $\mathbf{y}_1, \mathbf{y}_2 \in \mathbf{Pf}^{\text{sm}}$, we have

$$\text{Ext}^{j+2}(\mathcal{O}_{S_{\mathbf{y}_1}}, \mathcal{O}_{S_{\mathbf{y}_2}}(-j-1)) = 0.$$

3.1. Let $K \subset V$ be a linear subspace. Define $S \subset \mathbf{G}$ to be the locus of two-planes $T \subset V$ (i.e., points $T \in \mathbf{G}$) which intersect K non-trivially. Note that this is precisely the Schubert cycle corresponding to the increasing sequence $(0, \dim K, 7)$. The obvious $\text{GL}(V)$ action permutes the Schubert cycles corresponding to various subspaces K of V .

3.2. Proposition. *Let $K \subset V$ be a linear subspace. Regard $\wedge^2 \text{Ann } K \subset \wedge^2 V^*$ as a set of linear equations on \mathbf{P} . Then the set of two-planes $T \subset V$ such that $T \cap K \neq \emptyset$ is precisely the set of closed points in \mathbf{G} cut out by $\wedge^2 \text{Ann } K$.*

Proof. We have the following sequence of equivalent statements:

- $T \cap K \neq \emptyset$;
- the image \bar{T} of T in V/K has dimension at most one;
- $\wedge^2 \bar{T} = 0$ in $\wedge^2 V/K$;
- for every $\bar{w} \in \wedge^2 (V/K)^*$ we have $\bar{w}(\wedge^2 \bar{T}) = 0$;
- for every $w \in \wedge^2 \text{Ann } K$ we have $w(\wedge^2 T) = 0$.

(Note that the image of the map $\wedge^2 (V/K)^* \rightarrow \wedge^2 V^*$ is precisely $\wedge^2 \text{Ann } K$.) □

3.3. The above proposition shows that S is a closed subset of \mathbf{G} , and we endow it with the reduced induced scheme structure.

3.4. From now on, assume that $\dim K = 3$. Because of the $\text{GL}(V)$ action, we can assume that in a basis e_1, \dots, e_7 of V , K is spanned by e_5, e_6, e_7 . Let x_1, \dots, x_7 be the dual basis of V^* . The subspace $\text{Ann } K \subset V^*$, is spanned by x_1, \dots, x_4 , and thus $\wedge^2 \text{Ann } K$ has basis

$$x_1 \wedge x_2, x_1 \wedge x_3, \dots, x_3 \wedge x_4.$$

3.5. Proposition. *The Schubert cycle S is cut out scheme-theoretically by $\wedge^2 \text{Ann } K$. It is a rational variety with rational Cohen-Macaulay singularities, of codimension three in \mathbf{G} .*

Proof. In the Zariski open subspace $\mathbf{U} \cong \mathbf{C}^{10} \subseteq \mathbf{G}$ in which points T are given by

$$T = \text{Span}(\mathbf{a}_{11}\mathbf{e}_1 + \dots + \mathbf{a}_{15}\mathbf{e}_5 + \mathbf{e}_6, \mathbf{a}_{21}\mathbf{e}_1 + \dots + \mathbf{a}_{25}\mathbf{e}_5 + \mathbf{e}_7),$$

the cycle S is characterized by the condition that the matrix $R = \{\mathbf{a}_{ij}\}$, $i = 1, 2$, $j = 1, \dots, 4$ has rank one. The equations from $\wedge^2 \text{Ann } K$ are spanned by $x_i \wedge x_j$, $1 \leq i < j \leq 4$, which are precisely the maximal minors of the matrix R . It is a well-known fact that this determinantal variety is reduced and irreducible, and it coincides with the product of \mathbf{C}^2 and the cone over the Segre embedding of $\mathbf{P}^1 \times \mathbf{P}^3$. Thus $S \cap \mathbf{U}$ is cut out by $\wedge^2 \text{Ann } K$ and is rational. It remains to observe that every point in \mathbf{G} can be mapped inside \mathbf{U} by an element of $\text{GL}(V)$ that fixes K .

The singularities of S are rational Cohen-Macaulay since $S \cap \mathbf{U}$ is a toric variety (also see Propositions 3.8 and 3.6). \square

3.6. Proposition. *Denote by T the rank two tautological bundle on \mathbf{G} , and let $A = \text{Ann } K$. Then there exists a resolution of \mathcal{O}_S on \mathbf{G} of the form:*

$$0 \rightarrow \wedge^4 A \otimes \text{Sym}^2(T)(-1) \rightarrow \wedge^3 A \otimes T(-1) \rightarrow \wedge^2 A \otimes \mathcal{O}_{\mathbf{G}}(-1) \rightarrow \mathcal{O}_{\mathbf{G}} \rightarrow \mathcal{O}_S \rightarrow 0.$$

Proof. In the local chart described in the proof of Proposition 3.5, the above resolution becomes the Eagon-Northcott complex resolving the locus of rank one matrices inside the space of 2×4 matrices

$$\begin{pmatrix} \mathbf{a}_{11} & \mathbf{a}_{12} & \mathbf{a}_{13} & \mathbf{a}_{14} \\ \mathbf{a}_{21} & \mathbf{a}_{22} & \mathbf{a}_{23} & \mathbf{a}_{24} \end{pmatrix}.$$

(For a reference on Eagon-Northcott complexes, see [2, Appendix A2].) \square

3.7. We are now interested in getting a better understanding of the geometry of S . Let \tilde{S} be the set of pairs (T_1, T_2) of linear subspaces of V , of dimensions 1 and 2, respectively, such that $T_1 \subseteq T_2 \cap K$. Since $T_1 \subset K$ is a one-dimensional linear subset, the first projection μ maps \tilde{S} to $\mathbf{P}(K)$. Similarly, the second projection π maps \tilde{S} to S . Thus we have the diagram

$$\begin{array}{ccc} \tilde{S} & \xrightarrow{\mu} & \mathbf{P}(K) \\ \pi \downarrow & & \\ S & & \end{array}$$

Giving a point $(T_1, T_2) \in \tilde{S}$ amounts to choosing a line $T_1 \subset K$, along with another line T_2/T_1 in the six-dimensional space V/T_1 . Expressing this globally, we see that \tilde{S} is the projectivization of the rank six vector bundle on $\mathbf{P}(K)$

$$(V \otimes \mathcal{O})/\mathcal{O}(-1).$$

3.8. Proposition. *The map π is a resolution of singularities of S . There is a short exact sequence of bundles on \tilde{S}*

$$0 \rightarrow \mu^* \mathcal{O}_{\mathbf{P}(K)}(-1) \rightarrow \pi^* T \rightarrow \mathcal{O}_\mu(-1) \rightarrow 0,$$

where T is the tautological rank two vector bundle on \mathbf{G} , restricted to S , and $\mathcal{O}_\mu(-1)$ is the tautological relative bundle of the projection μ . We have

$$\pi^* \mathcal{O}_S(-1) = \mu^* \mathcal{O}_{\mathbf{P}(K)}(-1) \otimes \mathcal{O}_\mu(-1).$$

Proof. The space \tilde{S} is smooth, and π is obviously birational, thus π is a resolution of singularities. The short exact sequence is nothing but a global way of expressing the fact that T_2 is an extension of T_1 and T_2/T_1 . The last equality follows from the fact that on \mathbf{G} we have

$$\wedge^2 T = \mathcal{O}_{\mathbf{G}}(-1).$$

□

3.9. Proposition. *For $3 \leq j \leq 5$, $0 \leq k \leq 2$, the bundles $\mathcal{O}(j-6)$ and $\text{Sym}^k(T^*)(-j)$ have no cohomology on S .*

Proof. Since the singularities of S are rational, we can work on \tilde{S} by replacing all the bundles by their pull-backs by π . Using Proposition 3.8 we see that T^* is an extension of $\mu^* \mathcal{O}_{\mathbf{P}(K)}(1)$ and $\mathcal{O}_\mu(1)$. Thus $\text{Sym}^2(T^*)$ has a filtration with associated graded pieces

$$\mu^* \mathcal{O}_{\mathbf{P}(K)}(2), \mu^* \mathcal{O}_{\mathbf{P}(K)}(1) \otimes \mathcal{O}_\mu(1), \mathcal{O}_\mu(2).$$

Using the computation of $\mathcal{O}(-1)$ from the same Proposition, it follows that all the bundles we are interested in have filtrations whose associated graded object is a sum of line bundles of the form

$$\mu^* \mathcal{O}_{\mathbf{P}(K)}(k) \otimes \mathcal{O}_\mu(l),$$

with $-6 \leq l \leq -1$. By the Leray spectral sequence, these line bundles have no cohomology. □

3.10. We are now ready to define the Schubert cycles $S_{\mathbf{y}}$, parametrized by the smooth points of \mathbf{Pf} . Let $\mathbf{y} \in \mathbf{Pf}$ be a smooth point, which is thought of as a two-form on V of rank four. Let $K_{\mathbf{y}}$ be the kernel of this form, a linear three-space in V . We define the locus $S_{\mathbf{y}} \subset \mathbf{G}$ to be the Schubert cycle S defined above, for $K = K_{\mathbf{y}}$.

3.11. Proposition. *There exists a subvariety \mathbf{S} of $\mathbf{G} \times \mathbf{Pf}^{\text{sm}}$, flat over \mathbf{Pf}^{sm} , whose fiber $S_{\mathbf{y}} \subset \mathbf{G}$ over any closed point $\mathbf{y} \in \mathbf{Pf}^{\text{sm}}$ is precisely $S_{\mathbf{y}}$.*

Proof. Fix a point $\mathbf{y}_0 \in \mathbf{Pf}^{\text{sm}}$, and let S_0 be the corresponding subvariety of \mathbf{G} . The group $\text{GL}(V)$ acts transitively on both \mathbf{G} and \mathbf{Pf}^{sm} . One can take \mathbf{S} to be the image of $\text{GL}(V) \times S_0$ under this action, with the reduced-induced scheme structure. The flatness statement follows from [3, Theorem 9.9], as all translates of S_0 have the same Hilbert polynomial in \mathbf{G} . □

3.12. Proposition. (G-vanishing). *Let S_1, S_2 be the Schubert varieties defined in (3.10), corresponding to linear three-dimensional subspaces K_1, K_2 of V , and let $\mathcal{I}_{S_1}, \mathcal{I}_{S_2}$ denote their ideal sheaves in \mathbf{G} . Then, for $0 \leq j \leq 5$, we have*

$$\mathrm{Ext}_{\mathbf{G}}^{\bullet}(\mathcal{I}_{S_1}, \mathcal{I}_{S_2}(-j-1)) = 0.$$

Proof. In view of Serre duality and the fact that $K_{\mathbf{G}} = \mathcal{O}_{\mathbf{G}}(-7)$, it suffices to show that for $j = 3, 4$, or 5 , we have

$$\mathrm{Ext}_{\mathbf{G}}^{\bullet}(\mathcal{I}_{S_1}, \mathcal{I}_{S_2}(-j-1)) = 0.$$

From the long exact sequence

$$0 \rightarrow \mathcal{I}_{S_2}(-j-1) \rightarrow \mathcal{O}_{\mathbf{G}}(-j-1) \rightarrow \mathcal{O}_{S_2}(-j-1) \rightarrow 0$$

we will get the desired vanishing if we prove that

$$\mathrm{Ext}_{\mathbf{G}}^{\bullet}(\mathcal{I}_{S_1}, \mathcal{O}_{\mathbf{G}}(-j-1)) = \mathrm{Ext}_{\mathbf{G}}^{\bullet}(\mathcal{I}_{S_1}, \mathcal{O}_{S_2}(-j-1)) = 0.$$

By Serre duality and standard facts about the cohomology of $\mathcal{O}_{\mathbf{G}}$, the first statement amounts to proving that

$$\mathrm{H}^{\bullet}(\mathbf{G}, \mathcal{I}_{S_1}(j-6)) = \mathrm{H}^{\bullet}(S_1, \mathcal{O}_{S_1}(j-6)) = 0,$$

which is part of Proposition 3.9.

On the other hand, the resolution from Proposition 3.6 together with the local-to-global spectral sequence will give us the second vanishing statement provided we argue that

$$\mathrm{H}^{\bullet}(S_2, \mathrm{Sym}^k(\mathrm{T}^*(-j))) = 0$$

for $0 \leq k \leq 2$. This is also part of Proposition 3.9. □

3.13. As pointed out by Kuznetsov, **G**-vanishing also follows from [7, Theorem 3.1], using the resolution obtained in Proposition 3.6.

4. A family of curves

In this section we define, for a point $y \in Y$, a curve C_y in X . (We abuse the notation slightly: C_y may not be reduced or irreducible, but it does have dimension one.) The family $\{\mathcal{I}_{C_y}\}_{y \in Y}$ of ideal sheaves of these curves is the orthogonal family which induces the equivalence of derived categories $\mathbf{D}_{\mathrm{coh}}^b(X) \cong \mathbf{D}_{\mathrm{coh}}^b(Y)$. We then argue that $\dim C_y = 1$ for every choice of $y \in Y$ (which is essential in proving that the family of C_y 's is flat), and that

$$y_1 \neq y_2 \implies \mathrm{Hom}_X(\mathcal{I}_{C_{y_1}}, \mathcal{I}_{C_{y_2}}) = 0,$$

which is later needed for the orthogonality of the family.

Most of our statements depend on the assumption that X and Y are smooth. Therefore, for the remainder of this paper we shall assume that a space $W \subset \wedge^2 V^*$ is chosen in such a way that Y (and therefore X) is smooth.

4.1. Lemma. *Let $W \subset \wedge^2 V^*$ be a seven-dimensional subspace, and assume that $Y = \mathbf{W} \cap \mathbf{P}f$ is smooth of dimension three. Then the kernels of any pair of points $w_1 \neq w_2$ in Y (thought of as two-forms on V) are distinct.*

Proof. Assume that $\text{Ker } w_1 = \text{Ker } w_2 = K$. All the linear combinations of w_1 and w_2 have K contained in their kernel, so we can think of them as a one-dimensional family of forms on the four-dimensional space V/K . At some point w in this family the rank will drop to two, as this is a codimension one condition. At w the space $\mathbf{P}f$ is singular, and therefore the regular cut Y is singular at w as well. \square

4.2. Lemma. *Assume that X is smooth of dimension three. Then the space of linear forms $w \in \mathbf{P}^*$ which vanish along all of X is precisely W .*

Proof. Another way to phrase this assertion is by saying that W is the kernel of the map

$$\rho : H^0(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(1)) \rightarrow H^0(X, \mathcal{O}_X(1)).$$

Since the space W is already contained in the kernel of ρ , it suffices to show that $\dim \text{Ker } \rho = 7$. Factor ρ into the composition

$$H^0(\mathbf{P}, \mathcal{O}_{\mathbf{P}}(1)) \rightarrow H^0(\mathbf{G}, \mathcal{O}_{\mathbf{G}}(1)) \rightarrow H^0(X, \mathcal{O}_X(1)),$$

where it is well-known that the first map is an isomorphism.

Consider the ideal sheaf \mathcal{I}_X of X in \mathbf{G} . Its twist by one has the Koszul resolution

$$0 \rightarrow \mathcal{O}_{\mathbf{G}}(-6) \rightarrow \cdots \rightarrow \mathcal{O}_{\mathbf{G}}(-1)^{21} \rightarrow \mathcal{O}_{\mathbf{G}}^7 \rightarrow \mathcal{I}_X(1) \rightarrow 0,$$

where the sheaves $\mathcal{O}_{\mathbf{G}}(-1), \dots, \mathcal{O}_{\mathbf{G}}(-6)$ are acyclic with no global sections. It follows that

$$H^0(\mathbf{G}, \mathcal{I}_X(1)) = H^0(\mathbf{G}, \mathcal{O}_{\mathbf{G}}^7) = \mathbf{C}^7.$$

From this and the short exact sequence

$$0 \rightarrow \mathcal{I}_X(1) \rightarrow \mathcal{O}_{\mathbf{G}}(1) \rightarrow \mathcal{O}_X(1) \rightarrow 0$$

it follows that $\dim \text{Ker } \rho = 7$. \square

4.3. Let \mathbf{C} be the scheme theoretic intersection of \mathbf{S} with $X \times Y$ inside $\mathbf{G} \times \mathbf{P}f$, regarded as a subscheme of $X \times Y$. The sheaf $\mathcal{I}_{\mathbf{C}}$, the ideal sheaf of \mathbf{C} in $X \times Y$, regarded as an object of $\mathbf{D}_{\text{coh}}^b(X \times Y)$, will be the kernel of the Fourier-Mukai transform $\mathbf{D}_{\text{coh}}^b(Y) \rightarrow \mathbf{D}_{\text{coh}}^b(X)$.

4.4. Proposition. *The fiber of \mathbf{C} over any closed point $y \in Y$ has dimension one. The scheme \mathbf{C} is flat over Y .*

Proof. By an appropriate version of Bertini's theorem, we can choose a sequence of cuts going from \mathbf{G} to X so that all the intermediate members are smooth. Lefschetz's theorem then gives $\text{Pic } X = \mathbf{Z}$, generated by the restriction of $\mathcal{O}_{\mathbf{G}}(1)$. For a point $y \in Y$, C_y is cut out from X by the same six linear equations that cut out S_y from \mathbf{G} , i.e., C_y is the result of intersecting X with the hyperplanes corresponding to the points of the six-dimensional linear space $L = \wedge^2 \text{Ann Ker } y$. Every form $w \in L$ contains $\text{Ker } y$ in its kernel.

Assume by contradiction that C_y contains a divisor D . We claim that the intersection $H \cap X$ is either D , or all of X , for every hyperplane H in L . Indeed, any non-trivial intersection

$H \cap X$ is the generator of $\text{Pic } X$, and as such it is a prime divisor on X . Therefore any such cut must equal D , since it already contains it.

Pick a point P in $X \setminus D$. The condition that H contain P is a single linear condition on L , thus there is a five-dimensional space R of forms in L that vanish at P . The above discussion shows that if a form in L vanishes at a point outside D , then it vanishes on all of X . Thus the forms in R vanish on all of X .

By Lemma 4.2, then, we have $R \subset W$. Any two linearly-independent forms of rank four in R yield two points in Y whose kernel (as forms) is K . By Lemma 4.1, this contradicts the assumption that Y is smooth. Therefore our assumption that C_y contains a divisor must be false.

The subvarieties $S_y \subset \mathbf{G}$ for $y \in Y$ are all isomorphic, and they have the same Hilbert polynomial (in fact, there is a transitive $\text{GL}(V)$ -action permuting them). If we let y, y_1, \dots, y_6 be a basis of W , then C_y is obtained from S_y by six cuts with hyperplanes corresponding to y_1, \dots, y_6 . Since C_y is a curve (codimension six in W), it follows that this is a regular sequence of cuts, and thus the Hilbert polynomial of C_y does not depend on $y \in Y$. Therefore the family $\{C_y\}_{y \in Y}$ is flat over Y , by [3, III.9.9]. \square

4.5. Proposition. *Let y be a point in Y . Then we have*

$$\text{Hom}_X(\mathcal{I}_{C_y}, \mathcal{I}_{C_y}) = \mathbf{C}.$$

Proof. From the short exact sequence

$$0 \rightarrow \mathcal{I}_{C_y} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_{C_y} \rightarrow 0$$

we get

$$0 \rightarrow \mathbf{C} \rightarrow \text{Hom}_X(\mathcal{I}_{C_y}, \mathcal{O}_X) \rightarrow \text{Ext}_X^1(\mathcal{O}_{C_y}, \mathcal{O}_X) = H^2(X, \mathcal{O}_{C_y})^* = 0.$$

Thus any non-zero homomorphism $\mathcal{I}_{C_y} \rightarrow \mathcal{O}_X$ is, up to a scalar multiple, the usual inclusion.

Let $f : \mathcal{I}_{C_y} \rightarrow \mathcal{I}_{C_y}$ be any non-zero homomorphism. Composing with the inclusion $\mathcal{I}_{C_y} \hookrightarrow \mathcal{O}_X$ we get a non-zero map $\mathcal{I}_{C_y} \rightarrow \mathcal{O}_X$ which, by the above reasoning, must be the usual inclusion of \mathcal{I}_{C_y} into \mathcal{O}_X . Therefore f is injective, and since \mathcal{I}_{C_y} is torsion-free, it follows that f is a multiple of the identity map. \square

5. A reduction to the Grassmannian

In this section we argue that checking the orthogonality of the family $\{\mathcal{I}_{C_y}\}_{y \in Y}$ can be reduced to the \mathbf{G} -vanishing statement (Proposition 3.12).

5.1. We assume that we have chosen W so that X and Y are smooth of dimension three. Let y_1 and y_2 be two distinct points in Y . Regarded as smooth points of the Pfaffian \mathbf{Pf} , y_1 and y_2 give rise to Schubert cycles S_1 and S_2 in \mathbf{G} as explained in (3.10). These are rational, integral, codimension three subschemes of \mathbf{G} , whose singularities are rational and Cohen-Macaulay.

Let C_1 and C_2 be the corresponding curves in X obtained by taking scheme-theoretic intersections of S_1 and S_2 with X , respectively, as defined in Section 4.

We recall the \mathbf{G} -vanishing statement of Proposition 3.12:

$$\text{Ext}_{\mathbf{G}}^{\bullet}(\mathcal{I}_{S_1}, \mathcal{I}_{S_2}(-j-1)) = 0 \text{ for } 0 \leq j \leq 5.$$

5.2. Notation. Throughout this section, if S is a subscheme of a scheme Z , we shall denote by \mathcal{I}_S the ideal sheaf of S , regarded as a coherent sheaf on Z . In cases where this notation may be ambiguous, we shall make sure to specify which ambient scheme we are referring to.

5.3. Proposition. For $y_1 \neq y_2$ we have

$$\mathrm{Ext}_X^\bullet(\mathcal{I}_{C_1}, \mathcal{I}_{C_2}) = 0.$$

5.4. Before we begin the proof of Proposition 5.3 we need several intermediate vanishing results.

5.5. Lemma. For $0 \leq j \leq 5$ we have

$$H^\bullet(\mathbf{G}, \mathcal{I}_{S_2}(-j)) = 0.$$

Proof. Consider the short exact sequence

$$0 \rightarrow \mathcal{I}_{S_2} \rightarrow \mathcal{O}_{\mathbf{G}} \rightarrow \mathcal{O}_{S_2} \rightarrow 0.$$

The cohomology groups

$$H^\bullet(\mathbf{G}, \mathcal{O}_{\mathbf{G}}(-j))$$

vanish for $1 \leq j \leq 6$, by Kodaira vanishing. Since S_2 is irreducible, rational, and with rational singularities, we conclude that

$$H^\bullet(\mathbf{G}, \mathcal{I}_{S_2}) = 0,$$

which is the $j = 0$ case. For $1 \leq j \leq 5$ we have

$$H^{\bullet+1}(\mathbf{G}, \mathcal{I}_{S_2}(-j)) = H^\bullet(S_2, \mathcal{O}_{S_2}(-j)),$$

which vanishes by Proposition 3.9. □

5.6. We now divide the proof of Proposition 5.3 into several steps, corresponding to decreasing subvarieties of \mathbf{G} obtained by successive cuts with hyperplanes of increasing codimension, corresponding to subspaces of \mathbf{W} of increasing dimension.

To begin, let H_1 be the hyperplane in \mathbf{P} corresponding to y_1 , and let Z_1 be the scheme-theoretic intersection of \mathbf{G} and H_1 . It is a hypersurface of \mathbf{G} whose singularities are locally isomorphic to the product of \mathbf{C}^2 and an ordinary double point of dimension seven. Consider L_1 and L_2 which are the intersection of S_1 and S_2 with H_1 , regarded as subschemes of Z_1 . While H_1 cuts down the dimension of S_2 by one, it already contains S_1 . (The fact that H_1 cuts down the dimension of S_2 follows from Proposition 4.4.) Thus L_1 and L_2 have codimensions two and three, respectively, in Z_1 .

5.7. Proposition. *\mathbf{G} -vanishing implies Z_1 -vanishing:*

$$\mathrm{Ext}_{Z_1}^\bullet(\mathcal{I}_{L_1}, \mathcal{I}_{L_2}(-j)) = 0 \text{ for } 0 \leq j \leq 5.$$

Proof. Let $r : Z_1 \rightarrow \mathbf{G}$ denote the natural embedding. Since L_2 is obtained by a transversal cut of S_2 by H_1 , we have $\mathcal{I}_{L_2} = r^* \mathcal{I}_{S_2}$. By an easy form of Grothendieck duality it follows that

$$\begin{aligned} \mathrm{Ext}_{Z_1}^\bullet(\mathcal{I}_{L_1}, \mathcal{I}_{L_2}(-j)) &= \mathrm{Ext}_{Z_1}^\bullet(\mathcal{I}_{L_1}, r^* \mathcal{I}_{S_2}(-j)) \\ &= \mathrm{Ext}_{\mathbf{G}}^\bullet(r_! \mathcal{I}_{L_1}, \mathcal{I}_{S_2}(-j)) \\ &= \mathrm{Ext}_{\mathbf{G}}^{\bullet+1}(r_* \mathcal{I}_{L_1}, \mathcal{I}_{S_2}(-j-1)), \end{aligned}$$

since the embedding $Z_1 \subset \mathbf{G}$ has relative dualizing complex $\mathcal{O}_{Z_1}(1)[-1]$.

Consider the short exact sequence of sheaves on \mathbf{G}

$$0 \rightarrow \mathcal{I}_{Z_1} = \mathcal{O}_{\mathbf{G}}(-1) \rightarrow \mathcal{I}_{S_1} \rightarrow r_* \mathcal{I}_{L_1} \rightarrow 0.$$

Writing down the long exact sequence of Ext 's we get the result from \mathbf{G} -vanishing and the vanishing of

$$\mathrm{Ext}_{\mathbf{G}}^\bullet(\mathcal{I}_{Z_1}, \mathcal{I}_{S_2}(-j-1)) = \mathrm{H}^\bullet(\mathbf{G}, \mathcal{I}_{S_2}(-j)),$$

which is Lemma 5.5. □

5.8. Lemma. *For $0 \leq j \leq 5$ we have*

$$\mathrm{H}^\bullet(Z_1, \mathcal{I}_{L_1}(-j)) = 0.$$

Proof. The proof is essentially the same as that of Lemma 5.5 and will be left to the reader. One needs to use the fact that Z_1 is rational with rational singularities and $\mathrm{K}_{Z_1} = \mathcal{O}(-6)$. □

5.9. We can now complete the proof of Proposition 5.3. Let H_2 be the hyperplane in \mathbf{P} corresponding to \mathfrak{y}_2 , and let $Z_{1,2} = \mathbf{G} \cap H_1 \cap H_2$. Let D_1, D_2 be the intersections of S_1 and S_2 with $Z_{1,2}$, regarded as subschemes of $Z_{1,2}$. Note that since $S_2 \subset H_2$, we have $D_2 = L_2$; both D_1 and D_2 are now codimension two in $Z_{1,2}$.

Let W_5 be any linear subspace of W which, together with the one-dimensional subspaces corresponding to \mathfrak{y}_1 and \mathfrak{y}_2 , spans W . Because of dimension reasons, X, C_1 , and C_2 are obtained from $Z_{1,2}, D_1$, and D_2 by five successive transversal cuts with five hyperplanes corresponding to a basis of W_5 .

The strategy of the proof is to look at the successive embeddings

$$X \xrightarrow{g} Z_{1,2} \xrightarrow{h} Z_1 \xrightarrow{r} \mathbf{G},$$

and to use the vanishing of Ext groups on each one to conclude the vanishing of other Ext groups on the previous one. We have already argued (Proposition 5.7) that \mathbf{G} -vanishing implies Z_1 -vanishing. The passage from Z_1 -vanishing to the required vanishing on X will be done by an appropriate resolution of $h_* g_* \mathcal{I}_{C_2}$ on Z_1 .

We have

$$\mathcal{I}_{C_1} = g^* \mathcal{I}_{D_1},$$

thus

$$\mathrm{Ext}_{\mathcal{X}}^\bullet(\mathcal{I}_{C_1}, \mathcal{I}_{C_2}) = \mathrm{Ext}_{\mathcal{X}}^\bullet(g^* \mathcal{I}_{D_1}, \mathcal{I}_{C_2}) = \mathrm{Ext}_{Z_{1,2}}^\bullet(\mathcal{I}_{D_1}, g_* \mathcal{I}_{C_2}),$$

and

$$g_* \mathcal{I}_{C_2} = \mathcal{I}_{D_2} \otimes_{Z_{1,2}} \mathcal{O}_X.$$

Because the intersection of D_2 and X inside $Z_{1,2}$ is transversal, there are no higher Tor's in the above tensor product. We conclude that $g_* \mathcal{I}_{C_2}$ is quasi-isomorphic to the complex

$$\begin{aligned} \mathcal{I}_{D_2} \otimes \mathrm{Koszul}^{Z_{1,2}}(W_5) = \\ (0 \rightarrow \mathcal{I}_{D_2}(-5) \otimes \wedge^5 W_5 \rightarrow \cdots \rightarrow \mathcal{I}_{D_2}(-1) \otimes W_5 \rightarrow \mathcal{I}_{D_2} \rightarrow 0). \end{aligned}$$

Now let us move on to the embedding $h : Z_{1,2} \rightarrow Z_1$. Again,

$$\mathrm{Ext}_{Z_{1,2}}^\bullet(\mathcal{I}_{D_1}, g_* \mathcal{I}_{C_2}) = \mathrm{Ext}_{Z_{1,2}}^\bullet(h^* \mathcal{I}_{L_1}, g_* \mathcal{I}_{C_2}) = \mathrm{Ext}_{Z_1}^\bullet(\mathcal{I}_{L_1}, h_* g_* \mathcal{I}_{C_2}).$$

We use the previous resolution of $g_* \mathcal{I}_{C_2}$ to compute a resolution of $h_* g_* \mathcal{I}_{C_2}$ on Z_1 . The main difference is that \mathcal{I}_{D_2} is not obtained from \mathcal{I}_{L_2} by a transversal intersection. In fact, L_2 is already contained in H_2 , so we have a short exact sequence on Z_1

$$0 \rightarrow \mathcal{I}_{Z_{1,2}} = \mathcal{O}_{Z_1}(-1) \rightarrow \mathcal{I}_{L_2} \rightarrow h_* \mathcal{I}_{D_1} \rightarrow 0,$$

and $h_* \mathcal{I}_{D_1}$ is quasi-isomorphic to the two-term complex

$$0 \rightarrow \mathcal{O}_{Z_1}(-1) \rightarrow \mathcal{I}_{L_2} \rightarrow 0.$$

Observe that the Koszul complex restricts from Z_1 to $Z_{1,2}$, because the five cuts from W_5 are transversal to Z_1 and $Z_{1,2}$. In other words,

$$h^* \mathrm{Koszul}^{Z_1}(W_5) = \mathrm{Koszul}^{Z_{1,2}}(W_5).$$

The projection formula for derived categories implies that

$$\begin{aligned} h_* g_* \mathcal{I}_{C_2} &= h_*(\mathrm{Koszul}^{Z_{1,2}}(W_5) \otimes \mathcal{I}_{D_1}) = h_*(h^* \mathrm{Koszul}^{Z_1}(W_5) \otimes \mathcal{I}_{D_1}) \\ &= \mathrm{Koszul}^{Z_1}(W_5) \otimes h_* \mathcal{I}_{D_1} = \mathrm{Koszul}^{Z_1}(W_5) \otimes (\mathcal{O}_{Z_1}(-1) \rightarrow \mathcal{I}_{L_2}). \end{aligned}$$

All operations above are to be understood as derived. The computations of derived tensor product, and derived pull-back are correct, as the Koszul complex is locally free. Since g and h are embeddings, the left hand side is just the one sheaf complex $h_* g_* \mathcal{I}_{C_2}$.

Thus $h_* g_* \mathcal{I}_{C_2}$ is quasi-isomorphic to the total complex of

$$\mathrm{Koszul}^{Z_1}(W_5) \otimes (\mathcal{O}_{Z_1}(-1) \rightarrow \mathcal{I}_{L_2}).$$

By the local-to-global spectral sequence, in order to conclude that

$$\mathrm{Ext}_{\mathcal{X}}^\bullet(\mathcal{I}_{C_1}, \mathcal{I}_{C_2}) = 0$$

it suffices to prove that, for $0 \leq j \leq 5$, we have

$$\mathrm{Ext}_{Z_1}^{\bullet}(\mathcal{I}_{L_1}, \mathcal{I}_{L_2}(-j)) = 0$$

and

$$\mathrm{Ext}_{Z_1}^{\bullet}(\mathcal{I}_{L_1}, \mathcal{O}_{Z_1}(-j-1)) = 0.$$

The first statement is precisely Z_1 -vanishing, which is implied by \mathbf{G} -vanishing by Proposition 5.7. The second statement follows by Serre duality: we have

$$\mathrm{Ext}_{Z_1}^{\bullet}(\mathcal{I}_{L_1}, \mathcal{O}_{Z_1}(-j-1)) = H^{\varrho-\bullet}(Z_1, \mathcal{I}_{L_1}(j-5))^*$$

which vanishes for $0 \leq j \leq 5$ by Lemma 5.8. \square

6. The derived equivalence

In this section we define the integral transform Φ , and we argue that it gives an equivalence of derived categories $\mathbf{D}_{\mathrm{coh}}^b(Y) \cong \mathbf{D}_{\mathrm{coh}}^b(X)$ by verifying the criterion of [1, Theorem 1.1].

6.1. Let \mathbf{C} be the dimension four subscheme of $X \times Y$ defined in (4.3). We take its ideal sheaf $\mathcal{I}_{\mathbf{C}}$ in $X \times Y$ as the kernel of an integral transform

$$\Phi : \mathbf{D}_{\mathrm{coh}}^b(Y) \rightarrow \mathbf{D}_{\mathrm{coh}}^b(X) \quad \Phi(\mathcal{E}) = \mathbf{R}\pi_{X,*}(\pi_Y^*(\mathcal{E}) \otimes^{\mathbf{L}} \mathcal{I}_{\mathbf{C}}).$$

(Here, π_X and π_Y are the projections from $X \times Y$ to X and Y , respectively.)

6.2. Theorem. *The functor Φ is a Fourier-Mukai transform, i.e., it is an equivalence of categories $\mathbf{D}_{\mathrm{coh}}^b(Y) \cong \mathbf{D}_{\mathrm{coh}}^b(X)$.*

Proof. Observe that since \mathbf{C} is flat over Y , $\Phi\mathcal{O}_{\mathbf{y}}$ is precisely the ideal sheaf $\mathcal{I}_{\mathbf{C}_{\mathbf{y}}}$ on X . Thus we can think of $\{\mathcal{I}_{\mathbf{C}_{\mathbf{y}}}\}_{\mathbf{y} \in Y}$ as a family of sheaves on X , parametrized by the points of Y . Moreover, this family satisfies the following properties:

1. $\mathrm{Ext}_X^i(\mathcal{I}_{\mathbf{C}_{\mathbf{y}_1}}, \mathcal{I}_{\mathbf{C}_{\mathbf{y}_2}}) = 0$ for any i , and any pair of distinct points $\mathbf{y}_1, \mathbf{y}_2$ in Y (Proposition 5.3);
2. $\mathcal{I}_{\mathbf{C}_{\mathbf{y}}}$ is a simple sheaf on X for $\mathbf{y} \in Y$ (Proposition 4.5);
3. X is Calabi-Yau, therefore $\mathcal{I}_{\mathbf{C}_{\mathbf{y}}} \otimes \omega_X \cong \mathcal{I}_{\mathbf{C}_{\mathbf{y}}}$.

Thus, in the terms of [1], the sheaf $\mathcal{I}_{\mathbf{C}}$ is strongly simple. Combining this with Property 3, Theorem 1.1 of [loc.cit.] shows that Φ is a Fourier-Mukai transform. \square

7. Connections with existing work

In this section we put the example in this paper in the context of Kuznetsov's Homological Projective Duality.

7.1. The derived equivalence that we obtain appears to be a particular case of Homological Projective Duality, as explained by Kuznetsov [6]. Let us recall the setting of this theory.

Let $X \subset \mathbf{P}(V)$ be a smooth projective variety, and assume that its derived category admits a semi-orthogonal decomposition of the form

$$\mathbf{D}_{\text{coh}}^b(X) = \langle \mathcal{A}_0, \mathcal{A}_1(1), \dots, \mathcal{A}_{i-1}(i-1) \rangle$$

where

$$0 \subseteq \mathcal{A}_0 \subseteq \mathcal{A}_1 \subseteq \dots \subseteq \mathcal{A}_{i-1}$$

are full subcategories of $\mathbf{D}_{\text{coh}}^b(X)$, and (i) denotes twisting by $\mathcal{O}_X(i)$. If H is a hyperplane in $\mathbf{P}(V)$, let X_H be the corresponding hyperplane section of X . Then it is easy to see that

$$\langle \mathcal{A}_1(1), \mathcal{A}_2(2), \dots, \mathcal{A}_{i-1}(i-1) \rangle$$

is a semiorthogonal collection in $\mathbf{D}_{\text{coh}}^b(X_H)$. In general there is no reason to expect this collection to generate $\mathbf{D}_{\text{coh}}^b(X_H)$. Let \mathcal{C}_H denote the orthogonal in $\mathbf{D}_{\text{coh}}^b(X_H)$ of the subcategory generated by the above collection.

We can think of $\{\mathcal{C}_H : H \in \mathbf{P}(V^*)\}$ as a family of triangulated categories, parametrized by $H \in \mathbf{P}(V^*)$. In certain cases we can find a smooth variety Y , together with a morphism $Y \rightarrow \mathbf{P}(V^*)$, such that $\{\mathbf{D}_{\text{coh}}^b(Y_H)\}_{H \in \mathbf{P}(V^*)}$ is essentially the family of \mathcal{C}_H 's (for details, see [6]). In this situation Y is called the Homological Projective Dual of X .

The main theorem of [6] is the statement that Y then admits a semiorthogonal decomposition of a similar form, and if L is a linear subspace of V^* , $L^\perp \subset V$ its annihilator, then the linear sections

$$X_L = X \times_{\mathbf{P}(V)} \mathbf{P}(L^\perp), \quad Y_L = Y \times_{\mathbf{P}(V^*)} \mathbf{P}(L)$$

have closely related derived categories $\mathbf{D}_{\text{coh}}^b(X_L)$, $\mathbf{D}_{\text{coh}}^b(Y_L)$. Explicitly, their categories will have several trivial components (for X_L , arising from the semiorthogonal decomposition of X , for Y_L from that of Y), as well as a component which is *the same* in X_L and Y_L . In particular, if the dimension of L is chosen properly, the trivial components will disappear, and we get

$$\mathbf{D}_{\text{coh}}^b(X_L) \cong \mathbf{D}_{\text{coh}}^b(Y_L).$$

Another important property of this setup is that X_L is smooth precisely when Y_L is, and thus Y is closely related to the classical projective dual of X .

7.2. In our setting, we have the smooth projective variety $\mathbf{G} \subset \mathbf{P}$, its projective dual \mathbf{Pf} , and linear cuts X and Y of them by dual linear spaces. The varieties X and Y are simultaneously smooth, and when they are, their derived categories are equivalent. It is very tempting, in this context, to conjecture the following.

7.3. Conjecture. *There exists a Lefschetz decomposition of the derived category of \mathbf{G} , and a smooth variety $\overline{\mathbf{Pf}} \rightarrow \mathbf{P}^*$ (possibly understood in an extended sense, for example a non-commutative scheme), mapping to \mathbf{Pf} , which is homologically projectively dual to \mathbf{G} with respect to the decomposition of $\mathbf{D}_{\text{coh}}^b(\mathbf{G})$. Furthermore, Theorem 0.3 is a direct application of the main result of [6].*

7.4. The current form of the conjecture was suggested by Kuznetsov, who pointed out that our choice of \mathbf{Pf} from an early version of the paper can not be the correct homologically projective dual. In fact, Kuznetsov claims to have constructed such a non-commutative homologically projective dual of $\mathbf{G}(2,7)$ in [8]. In addition, he expects to be able to extend this result to $\mathbf{G}(2,2n+1)$ for $n > 3$ as well.

References

- [1] Bridgeland, T., Equivalences of triangulated categories and Fourier-Mukai transforms, Bull. London Math. Soc. 31 (1999), no. 1, 25–34, (also preprint, [math.AG/9809114](#))
- [2] Eisenbud, D., *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics Vol. 150, Springer-Verlag (1995)
- [3] Hartshorne, R., *Algebraic Geometry*, Graduate Texts in Mathematics Vol. 52, Springer-Verlag (1977)
- [4] Hori, K., Tong, D., Aspects of non-abelian gauge dynamics in two-dimensional $\mathcal{N} = (2,2)$ theories, preprint, [hep-th/0609032](#)
- [5] Kollár, J., Mori, S., *Birational Geometry of Algebraic Varieties*, Cambridge Tracts in Mathematics Vol. 134, Cambridge University Press (1998)
- [6] Kuznetsov, A., Homological Projective Duality, preprint, [math.AG/0507292](#)
- [7] Kuznetsov, A., Exceptional collections for Grassmannians of isotropic lines, preprint, [math.AG/0512013](#)
- [8] Kuznetsov, A., Lefschetz exceptional collections and categorical resolutions of singularities, preprint, [math.AG/0609240](#)
- [9] Rødland, E., A., The Pfaffian Calabi-Yau, its Mirror, and their link to the Grassmannian $\mathbf{G}(2,7)$, *Compositio Math.* 122 (2000), no. 2, 135–149, (also preprint, [math.AG/9801092](#))