

MATH 844: HOMEWORK 1, DUE FEB 2

1. Suppose that (a, b) is a rational point on Reichardt's curve $x^4 - 17 = 2y^2$.

(a) Show that $a = A/C, b = B/C^2$, where $A, B, C \in \mathbf{Z}$, $\gcd(A, B) = \gcd(A, C) = \gcd(B, C) = 1$.

(b) Rewriting the equation as $(5A^2 + 17C^2)^2 - (4B)^2 = 17(A^2 + 5C^2)^2$, show that there are $U, V \in \mathbf{Z}$ such that

$$5A^2 + 17C^2 \pm 4B = 17U^2, 5A^2 + 17C^2 \mp 4B = V^2, A^2 + 5C^2 = UV$$

or

$$5A^2 + 17C^2 \pm 4B = 34U^2, 5A^2 + 17C^2 \mp 4B = 2V^2, A^2 + 5C^2 = 2UV$$

(c) Hence show that (eliminating B) either

$$10A^2 + 34C^2 = 17U^2 + V^2, A^2 + 5C^2 = UV$$

or

$$5A^2 + 17C^2 = 17U^2 + V^2, A^2 + 5C^2 = 2UV$$

(d) Show that neither possibility has a solution modulo 17, and so deduce that Reichardt's equation has no rational solutions.

(e) Show, however, that Reichardt's equation has a real solution and a solution modulo p for each prime number $p \equiv 3 \pmod{4}$. (It's possible but harder to show this for all primes p .)