

# A CONJECTURE OF EVANS ON SUMS OF KLOOSTERMAN SUMS

EVAN P. DUMMIT, ADAM W. GOLDBERG, AND ALEXANDER R. PERRY

ABSTRACT. In a recent paper [5], Evans relates twisted Kloosterman sheaf sums to Gaussian hypergeometric functions, and he formulates a number of conjectures relating certain twisted Kloosterman sheaf sums to the coefficients of modular forms. Here we prove one of his conjectures for a fourth order twisted Kloosterman sheaf sum  $T_n$  of the quadratic character on  $\mathbf{F}_p^\times$ . In the course of the proof we develop reductions for twisted moments of Kloosterman sums, and apply these in the end to derive a congruence relation for  $T_n$  with generalized Apéry numbers.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

Fix an odd prime  $p$ , and let  $\mathbf{F}_p$  be the finite field with  $p$  elements. For any multiplicative character  $\chi$  on  $\mathbf{F}_p^\times$ , we extend  $\chi$  to a function on  $\mathbf{F}_p$  by setting  $\chi(0) = 0$ . Let  $\varepsilon$  and  $\phi$  be the trivial and quadratic characters on  $\mathbf{F}_p^\times$ . Also let  $\psi : \mathbf{F}_p \rightarrow \mathbf{C}$  be the additive character  $x \mapsto e^{2\pi ix/p}$ . We shall write  $\bar{x}$  for the multiplicative inverse of  $x \in \mathbf{F}_p^\times$ . For  $a \in \mathbf{F}_p^\times$ , the classical *Kloosterman sum* is

$$(1.1) \quad K(a) := \sum_{x \in \mathbf{F}_p^\times} \psi(x + a\bar{x}).$$

The  $n$ th twisted *Kloosterman sheaf sum* of  $\phi$  is

$$(1.2) \quad T_n := \sum_{a \in \mathbf{F}_p^\times} \phi(a)(g(a)^n + g(a)^{n-1}h(a) + g(a)^{n-2}h(a)^2 + \cdots + g(a)h(a)^{n-1} + h(a)^n),$$

where  $g(a)$  and  $h(a)$  are the roots of the polynomial

$$(1.3) \quad X^2 + K(a)X + p.$$

We study a conjecture of Evans (see below) that relates  $T_4$  to the coefficient  $a(p)$  of the unique normalized newform in  $S_4(\Gamma_0(8))$ , which is given by

$$(1.4) \quad \sum_{n=1}^{\infty} a(n)q^n := q \prod_{n=1}^{\infty} (1 - q^{2n})^4 (1 - q^{4n})^4 = q - 4q^3 - 2q^5 + 24q^7 - 11q^9 - \cdots.$$

To do so, we employ Gaussian hypergeometric series. For multiplicative characters  $A$  and  $B$  on  $\mathbf{F}_p$ , define the normalized Jacobi sum

$$\binom{A}{B} = \frac{B(-1)}{p} J(A, \bar{B}) = \frac{B(-1)}{p} \sum_{x \in \mathbf{F}_p} A(x)\bar{B}(1-x).$$

For multiplicative characters  $A_0, \dots, A_n$  and  $B_1, \dots, B_n$  on  $\mathbf{F}_p$ , Greene [7] defined the *Gaussian hypergeometric series over  $\mathbf{F}_p$*  by

$${}_{n+1}F_n \left( \begin{matrix} A_0, & A_1, & \dots & A_n \\ & B_1, & \dots & B_n \end{matrix} \middle| x \right) = \frac{p}{p-1} \sum_{\chi} \binom{A_0\chi}{\chi} \binom{A_1\chi}{B_1\chi} \cdots \binom{A_n\chi}{B_n\chi} \chi(x),$$

where the sum is over all multiplicative characters  $\chi$  on  $\mathbf{F}_p$  (see also [11]). We will be interested in the case where the top parameters are the quadratic character  $\phi$  and the bottom ones the trivial character  $\varepsilon$ ; for brevity we write this function as

$$(1.5) \quad {}_{n+1}F_n(x) = {}_{n+1}F_n \left( \begin{matrix} \phi, & \phi, & \dots & \phi \\ & \varepsilon, & \dots & \varepsilon \end{matrix} \middle| x \right) = \frac{p}{p-1} \sum_{\chi} \binom{\phi\chi}{\chi}^{n+1} \chi(x).$$

In [2] and [3], Ahlgren and Ono prove  $p^3 {}_4F_3(1) = -a(p) - p$  by showing that the coefficients  $a(p)$  and the hypergeometric values  $p^3 {}_4F_3(1)$  count points on the ‘‘Calabi-Yau’’ variety

$$\{(x, y, z, w) \in (\mathbf{F}_p^\times)^4 \mid x + \frac{1}{x} + y + \frac{1}{y} + z + \frac{1}{z} + w + \frac{1}{w} = 0\}.$$

With this result, Evans’ Conjecture can be stated as follows.

**Conjecture** (Evans, [5]). *If  $p$  is an odd prime, then*

$$\frac{T_4}{p} = p^3 {}_4F_3(1) + p.$$

We prove the following theorem:

**Theorem 1.1.** *Evans’ Conjecture is true.*

Ahlgren and Ono’s work in [3] was motivated by Beukers’ Conjecture relating the coefficient  $a(p)$  to the Apéry number<sup>1</sup>  $A\left(\frac{p-1}{2}\right)$  where

$$(1.6) \quad A(n) := \sum_{j=0}^n \binom{n+j}{j}^2 \binom{n}{j}^2.$$

Beukers’ conjecture says that

$$(1.7) \quad A\left(\frac{p-1}{2}\right) \equiv a(p) \pmod{p^2}.$$

Ahlgren and Ono proved this conjecture using combinatorial properties of  ${}_4F_3(1)$  and the relation  $p^3 {}_4F_3(1) = -a(p) - p$  [3]. Since Evans’ conjecture relates  ${}_4F_3(1)$  to  $T_4$ , one expects that the  $T_n$  satisfy Beukers-type congruences. We derive a hypergeometric expression for the sum  $T_n$  and apply work of Ahlgren and Ono on hypergeometric congruences to prove the following.

**Theorem 1.2.** *For  $\lambda \in \mathbf{F}_p$ , define the generalized Apéry number by*

$$(1.8) \quad A(p, \lambda) := \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j}^2 \binom{\frac{p-1}{2} + j}{j} \lambda^{pj}$$

<sup>1</sup>Apéry used [4] the numbers  $A(n)$  in his famous proof that  $\zeta(3)$  is irrational.

for  $\lambda \in \mathbf{F}_p^\times$  and  $A(p, 0) = 1$ . Then, if  $n$  is a positive integer,

$$(1.9) \quad T_{n+3} \equiv (-1)^{n+1} p \sum_{x_1, \dots, x_n \in \mathbf{F}_p^\times} A \left( p, \frac{(x_1 + \dots + x_n + 1)(\bar{x}_1 + \dots + \bar{x}_n + 1)}{4} \right) \pmod{p^2}.$$

We briefly outline the structure of the paper. In the first section we provide background information on Gaussian hypergeometric functions. Next we prove reductions for  $T_n$ , including a purely hypergeometric expression as a sum of  ${}_3F_2$ 's; we then apply a finite field analogue of Clausen's theorem to the specific  $n = 4$  case. To complete the proof of Theorem 1.1, we express both  $T_4$  and  $p^3 {}_4F_3(1)$  as a "trace" of squares of  ${}_2F_1$ 's. In the last section we prove Theorem 1.2.

## 2. PRELIMINARIES ON GAUSSIAN HYPERGEOMETRIC FUNCTIONS

We state here some basic results on Gaussian hypergeometric functions. Any Gaussian hypergeometric function can be expressed as a "Diophantine" sum over  $\mathbf{F}_p$  [7]. We recall that for  $x \neq 0$ ,

$$(2.1) \quad {}_2F_1(x) = \frac{\phi(-1)}{p} \sum_{y \in \mathbf{F}_p^\times} \phi(y)\phi(1-y)\phi(1-xy),$$

$$(2.2) \quad {}_3F_2(x) = \frac{1}{p^2} \sum_{y, z \in \mathbf{F}_p^\times} \phi(y)\phi(1-y)\phi(z)\phi(1-z)\phi(1-xyz).$$

**Note.** Note that  ${}_3F_2(0) = {}_2F_1(0) = 0$  according to Greene's conventions given in the introduction; however, it is also natural to define  ${}_3F_2(0)$  and  ${}_2F_1(0)$  by the right hand sides of the equalities above. We will take this convention for the rest of the paper because it simplifies the statements of Theorems 3.1 and 1.2.

Like their classical counterparts, Gaussian hypergeometric functions satisfy many transformation laws. In this paper we need the following two transformations, which are applications of Theorems 4.2 and 4.4 of [7].

**Lemma 2.1.** For  $x \neq 0, 1$ , we have

$$(2.3) \quad {}_2F_1(x) = \phi(-1) {}_2F_1(1-x),$$

$$(2.4) \quad {}_2F_1(x) = \phi(x) {}_2F_1(1/x).$$

In what follows, we will need to express a  ${}_3F_2$  as a square of a  ${}_2F_1$ . To do so, we use a special case of a formula provided by Evans and Greene in Theorem 1.7 of [6].

**Lemma 2.2.** If  $t \neq 0, \pm 1$ , then

$${}_3F_2 \left( \frac{1}{1-t^2} \right) = \phi(t^2 - 1) \left( -\frac{1}{p} + {}_2F_1 \left( \frac{1-t}{2} \right)^2 \right).$$

The following expression for  $p^3 {}_4F_3(1)$  as a trace of squares of  ${}_2F_1$ 's is proven in Lemma 2.2 of [3].

**Lemma 2.3.** *We have that*

$$p^3 {}_4F_3(1) = p^2 \sum_{x \in \mathbf{F}_p^\times} \phi(x) {}_2F_1(x)^2.$$

Finally, we will need two special evaluations of hypergeometric functions from Corollary 11.12 and Theorem 11.14 of [11].

**Lemma 2.4.** *We have that*

$$(2.5) \quad {}_2F_1(-1) = \begin{cases} 0 & \text{if } p \equiv 3 \pmod{4}, \\ \frac{2x(-1)^{\frac{x+y+1}{2}}}{p} & \text{if } p \equiv 1 \pmod{4}, x^2 + y^2 = p, \text{ and } x \text{ odd}, \end{cases}$$

$$(2.6) \quad {}_3F_2(1) = \begin{cases} 0 & \text{if } p \equiv 3 \pmod{4}, \\ \frac{4x^2 - 2p}{p^2} & \text{if } p \equiv 1 \pmod{4}, x^2 + y^2 = p, \text{ and } x \text{ odd}. \end{cases}$$

### 3. REDUCTIONS OF TWISTED KLOOSTERMAN SUMS

Using the relations  $g(a) + h(a) = -K(a)$  and  $g(a)h(a) = p$ , which follow from (1.3), the polynomial  $g(a)^n + g(a)^{n-1}h(a) + \cdots + g(a)h(a)^{n-1} + h(a)^n$  can be written as a polynomial in  $K(a)$  with coefficients in  $\mathbf{Z}$ . Therefore  $T_n$  can be written as a linear combination of the twisted Kloosterman moments

$$(3.1) \quad S(m) = \sum_{a \in \mathbf{F}_p^\times} \phi(a) K(a)^m.$$

In this section we give general reductions for such exponential sums, relate them to hypergeometric functions, and specialize to the case  $T_4$ .

**3.1. Reductions of  $T_n$ .** The next proposition reduces the exponential sum  $S(m) = \sum_{a \in \mathbf{F}_p^\times} \phi(a) K(a)^m$  to a symmetric sum of quadratic characters.

**Proposition 3.1.** *For an integer  $m \geq 1$ , we have that*

$$S(m+1) = p\phi(-1) \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} \phi(x_1 + \cdots + x_m + 1) \phi(\overline{x_1} + \cdots + \overline{x_m} + 1).$$

*Proof.* By definition, we have that

$$\begin{aligned} \sum_{a \in \mathbf{F}_p^\times} \phi(a) K(a)^{m+1} &= \sum_{a \in \mathbf{F}_p^\times} \phi(a) \sum_{x_1, \dots, x_m, y \in \mathbf{F}_p^\times} \psi(x_1 + \cdots + x_m + y + a(\overline{x_1} + \cdots + \overline{x_m} + \overline{y})) \\ &= \sum_{x_1, \dots, x_m, y \in \mathbf{F}_p^\times} \psi(x_1 + \cdots + x_m + y) \sum_{a \in \mathbf{F}_p^\times} \phi(a) \psi(a(\overline{x_1} + \cdots + \overline{x_m} + \overline{y})). \end{aligned}$$

If  $\overline{x_1} + \cdots + \overline{x_m} + \overline{y} = 0$  then the sum on  $a$  vanishes, so we may multiply by

$$\phi(\overline{x_1} + \cdots + \overline{x_m} + \overline{y})^2 = \begin{cases} 0 & \text{if } \overline{x_1} + \cdots + \overline{x_m} + \overline{y} = 0 \\ 1 & \text{otherwise} \end{cases}$$

inside the sum on  $x_1, \dots, x_m, y$ . Thus we have

$$\begin{aligned} \sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)^{m+1} &= \sum_{x_1, \dots, x_m, y \in \mathbf{F}_p^\times} \phi(\sum_i \bar{x}_i + \bar{y})\psi(\sum_i x_i + y) \sum_{a \in \mathbf{F}_p^\times} \phi(a(\sum_i \bar{x}_i + \bar{y}))\psi(a(\sum_i \bar{x}_i + \bar{y})) \\ &= G(\phi) \sum_{x_1, \dots, x_m, y \in \mathbf{F}_p^\times} \phi(\sum_i \bar{x}_i + \bar{y})\psi(\sum_i x_i + y), \end{aligned}$$

where  $G(\phi) = \sum_{a \in \mathbf{F}_p^\times} \phi(a)\psi(a)$  is the usual Gauss sum. Making the change of variables  $(x_1, \dots, x_m, y) \mapsto (x_1y, x_2y, \dots, x_my, y)$ , we have

$$\sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)^{m+1} = G(\phi) \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} \phi(\sum_i \bar{x}_i + 1) \sum_{y \in \mathbf{F}_p^\times} \phi(y)\psi(y(\sum_i x_i + 1)).$$

As before, we can multiply by  $\phi(\sum_i x_i + 1)^2$  to get

$$\begin{aligned} \sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)^{m+1} &= G(\phi) \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} \phi(\sum_i \bar{x}_i + 1)\phi(\sum_i x_i + 1) \sum_{y \in \mathbf{F}_p^\times} \phi(y(\sum_i x_i + 1))\psi(y(\sum_i x_i + 1)) \\ &= G(\phi)^2 \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} \phi(\sum_i \bar{x}_i + 1)\phi(\sum_i x_i + 1). \end{aligned}$$

Since  $G(\phi)^2 = p\phi(-1)$  by Proposition 6.3.2 of [8], this finishes the proof.  $\square$

**Remark.** Note in particular that  $S(2) = -p$ . Also, the sum  $S(1) = \sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)$  is not covered by the proposition, but it is easy to evaluate  $S(1) = G(\phi)^2 = p\phi(-1)$  directly.

The form of the character sum in the proposition motivates the study of the auxiliary function

$$(3.2) \quad Q(a, b) = \sum_{x, y \in \mathbf{F}_p^\times} \phi(x + y + a)\phi(\bar{x} + \bar{y} + b)$$

for  $a, b \in \mathbf{F}_p$ . The function  $Q(a, b)$  serves as the connection between twisted Kloosterman sheaf sums and hypergeometric functions.

**Proposition 3.2.** *If  $a, b \in \mathbf{F}_p$  are not both zero, then  $Q(a, b) = p^2\phi(-1)_3F_2\left(\frac{ab}{4}\right) + p\phi(ab)$ . Additionally,  $Q(0, 0) = -(p-1)\phi(-1)$ .*

*Proof.* First, we may assume that  $a \neq 0$  by interchanging  $a$  and  $b$  (if necessary) because the sum is clearly symmetric in  $a$  and  $b$ . Now, for  $c \in \mathbf{F}_p^\times, d \in \mathbf{F}_p$ , we note the well-known formula

$$\sum_{x \in \mathbf{F}_p} \phi(cx^2 + dx + e) = \begin{cases} (p-1)\phi(c) & \text{if } d^2 - 4ce = 0, \\ -\phi(c) & \text{if } d^2 - 4ce \neq 0. \end{cases}$$

Applying the change of variables  $(x + y, xy) \mapsto (s, r)$  yields

$$\begin{aligned} Q(a, b) &= \sum_{s \in \mathbf{F}_p, r \in \mathbf{F}_p^\times} \phi(s + a)\phi(s/r + b)[1 + \phi(s^2 - 4r)] \\ &= \sum_{s \in \mathbf{F}_p, r \in \mathbf{F}_p^\times} \phi(s + a)\phi(s/r + b) + \sum_{r \in \mathbf{F}_p} \phi(a)\phi(b)\phi(-4r) + \sum_{r, s \in \mathbf{F}_p^\times} \phi(s + a)\phi(s/r + b)\phi(s^2 - 4r). \end{aligned}$$

The first sum equals  $p\phi(ab)$  by the note above, and the second sum clearly vanishes. Substituting  $(r, s) \mapsto (ra^2s^2/4, -as)$  in the third sum yields

$$\begin{aligned} \sum_{r,s \in \mathbf{F}_p^\times} \phi(r)\phi(s+a)\phi(s+br)\phi(s^2-4r) &= \sum_{r,s \in \mathbf{F}_p^\times} \phi(r)\phi(-as+a)\phi(-as+bra^2s^2/4)\phi(1-r) \\ &= \phi(-1) \sum_{r,s \in \mathbf{F}_p^\times} \phi(s)\phi(r)\phi(1-s)\phi(1-absr/4)\phi(1-r) \\ &= \phi(-1)p^2 {}_3F_2\left(\frac{ab}{4}\right). \end{aligned}$$

For the last statement of the proposition, we have

$$Q(0,0) = \sum_{x,y \in \mathbf{F}_p^\times} \phi(x+y)^2\phi(xy) = \sum_{x \in \mathbf{F}_p^\times} -\phi(-x^2) = (p-1)\phi(-1).$$

**Remark.** *The specific evaluation of  $Q(a,b)$  in the case  $a = b = 1$  is the subject of two notable early papers. The Lehmers gave the first evaluation of the sum, directly rewriting  $Q(1,1)$  as a Jacobsthal sum via a complicated change of variables [9]. A later paper by Mordell revisits their method, and finds a simpler path that also eventually leads to Jacobsthal sums [10]. The more general method here also allows for direct evaluation of  $Q(1,1) = \phi(-1)p^2 {}_3F_2(1/4) + p$ , as  ${}_3F_2(1/4)$  has an explicit evaluation as given in Theorem 11.17 of [11].* □

Combining Propositions 3.1 and 3.2 gives a development for  $T_n$  as sums of  ${}_3F_2$ 's. To state this precisely, we define for positive integers  $m$  the sum

$$(3.3) \quad G(m) := \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} {}_3F_2\left(\frac{(x_1 + x_2 + \dots + x_m + 1)(\overline{x_1} + \dots + \overline{x_m} + 1)}{4}\right).$$

**Theorem 3.1.** *For  $m \geq 1$ , there is a recurrence*

$$(3.4) \quad S(m+3) = p^3G(m) + pS(m+1) - p^2C(m),$$

where  $C(m)$  is the number of simultaneous solutions to the congruences  $x_1 + \dots + x_m = 0$  and  $\overline{x_1} + \dots + \overline{x_m} = 0$ .

*Proof.* Applying Propositions 3.1 and 3.2, we have

$$\begin{aligned} S(m+3) &= p\phi(-1) \sum_{x_1, \dots, x_{m+2} \in \mathbf{F}_p^\times} \phi(x_1 + \dots + x_{m+2} + 1)\phi(\overline{x_1} + \dots + \overline{x_{m+2}} + 1) \\ &= p\phi(-1) \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} Q(x_1 + \dots + x_m + 1, \overline{x_1} + \dots + \overline{x_{m+2}} + 1) \\ &= p^3G(m) + p^2\phi(-1) \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} \phi((x_1 + \dots + x_m + 1)(\overline{x_1} + \dots + \overline{x_m} + 1)) \\ &\quad + p\phi(-1)(Q(0,0) - p^2\phi(-1){}_3F_2(0))C(m) \\ &= p^3G(m) + pS(m+1) - p^2C(m) \end{aligned}$$

where the last term is a correction stemming from the fact that  $Q(0, 0) - p^2\phi(-1)_3F_2(0) = -p\phi(-1)$ , and this correction must be made exactly  $C(m)$  times.  $\square$

**Remark.** *This result allows us to express  $T_n$  as a linear combination of  ${}_3F_2$ 's,  $S(1)$ ,  $S(2)$ ,  $S(3)$  and the constants  $C(m)$ .*

**3.2. Simplifications of  $T_4$ .** A calculation using the relations  $g(a) + h(a) = -K(a)$  and  $g(a)h(a) = p$  shows that

$$g(a)^4 + g(a)^3h(a) + g(a)^2h(a)^2 + g(a)h(a)^3 + h(a)^4 = K(a)^4 - 3pK(a)^2 + p^2.$$

Therefore we have that

$$(3.5) \quad T_4 = \sum_{a \in \mathbf{F}_p^\times} \phi(a)(K(a)^4 - 3pK(a)^2 + p^2) = \sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)^4 - 3p \sum_{a \in \mathbf{F}_p^\times} \phi(a)K(a)^2.$$

We can now apply the results from the previous section to reduce  $T_4$ . To ease notation, we define a function  $F : \mathbf{F}_p^\times \rightarrow \mathbf{C}$  by

$$(3.6) \quad F(a) := Q(a + 1, \bar{a} + 1).$$

**Lemma 3.1.** *We have*

$$T_4 = p\phi(-1) \sum_{x \in \mathbf{F}_p^\times} F(x) + 3p^2.$$

*Proof.* Write  $T_4 = S(4) - 3pS(2)$  as in (3.5) and apply Proposition 3.1.  $\square$

**Lemma 3.2.** *For  $a \neq 0, \pm 1$ , we have that*

$$F(a) = p^2\phi(a)_2F_1(-a)^2.$$

*Proof.* By Proposition 3.2, for  $a \neq 0, -1$ , we have that

$$(3.7) \quad F(a) = p\phi(a) + p^2\phi(-1)_3F_2\left(\frac{(a+1)^2}{4a}\right).$$

Taking  $t = \frac{a-1}{a+1}$  in Lemma 2.2, we get

$$p^2\phi(-1)_3F_2\left(\frac{(a+1)^2}{4a}\right) = \phi(a) \left(-p + p^2_2F_1\left(\frac{1}{1+a}\right)^2\right),$$

and so  $F(a) = p^2\phi(a)_2F_1(1/(1+a))^2$ . Applying the two transformations in Lemma 2.1 finishes the proof.  $\square$

We will need two special evaluations of  $F$  at  $\pm 1$ , the points excluded from the previous lemma.

**Lemma 3.3.** *We have that*

$$(3.8) \quad \phi(-1)F(1) = p^2\phi(-1)_2F_1(-1)^2 - p$$

$$(3.9) \quad \phi(-1)F(-1) = p^2_2F_1(1)^2 - p$$

*Proof.* By Proposition 3.2 we have  $F(1) = p + p^2\phi(-1)_3F_2(1)$ , and so the first identity follows directly from Lemma 2.4. Note that

$$\begin{aligned}\phi(-1)F(-1) &= \phi(-1) \sum_{x,y \in \mathbf{F}_p^\times} \phi(x+y)\phi(\bar{x} + \bar{y}) = \phi(-1) \sum_{x,y \in \mathbf{F}_p^\times} \phi(xy)\phi(x+y)^2 \\ &= -\phi(-1) \sum_{x+y=0} \phi(xy) = -(p-1),\end{aligned}$$

and the Diophantine representation in (2.1) for  ${}_2F_1$  implies that  $p^2{}_2F_1(1)^2 = 1$ . This proves (3.9).  $\square$

#### 4. PROOF OF THEOREM 1.1

By successively applying Lemmas 3.1, 3.2, and 3.3, we have

$$\begin{aligned}\frac{T_4}{p} &= 3p + \phi(-1) \sum_{x \in \mathbf{F}_p^\times} F(x) \\ &= 3p + \phi(-1)F(1) + \phi(-1)F(-1) + p^2 \sum_{x \in \mathbf{F}_p^\times \setminus \{\pm 1\}} \phi(-x)_2F_1(-x)^2 \\ &= 3p + (\phi(-1)F(1) - p^2\phi(-1)_2F_1(-1)^2) + (\phi(-1)F(-1) - p^2{}_2F_1(1)^2) + p^2 \sum_{x \in \mathbf{F}_p^\times} \phi(x)_2F_1(x)^2 \\ &= p + p^2 \sum_{x \in \mathbf{F}_p^\times} \phi(x)_2F_1(x)^2.\end{aligned}$$

Applying Lemma 2.3 yields

$$\frac{T_4}{p} = p^3{}_4F_3(1) + p.$$

#### 5. APÉRY-TYPE CONGRUENCES

For the proof of Theorem 1.2, we recall a result of Ahlgren [1].

**Theorem 5.1** (Ahlgren). *If  $p$  is an odd prime and  $1 \leq \lambda \leq p-1$ , then define*

$$\begin{aligned}A(p, \lambda) &:= \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j}^2 \binom{\frac{p-1}{2} + j}{j} \lambda^{pj}, \\ B(p, \lambda) &:= \sum_{j=0}^{\frac{p-1}{2}} \binom{\frac{p-1}{2}}{j}^2 \binom{\frac{p-1}{2} + j}{j} \lambda^j \cdot \left\{ 1 + \frac{1}{2} \sum_{i=\frac{p+1}{2}}^{\frac{p-1}{2}+j} \frac{1}{i} + 3j \sum_{i=1+j}^{\frac{p-1}{2}+j} \frac{1}{i} \right\}.\end{aligned}$$

Then we have

$$(5.1) \quad p^2{}_3F_2(\lambda) \equiv A(p, \lambda) + pB(p, \lambda) \pmod{p^2}.$$

**Remark.** *If we define  $A(p, 0) = 1$  and  $B(p, 0) = 0$  then the results self-evidently hold also for  $\lambda = 0$ . We will use this definition in our proof below.*

*Proof of Theorem 1.2.* By Proposition 3.1,  $pS(m) \equiv 0 \pmod{p^2}$ . Therefore, Theorems 3.1 and 5.1 together show

$$\begin{aligned} S(m+3) &\equiv \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} p^3 {}_3F_2 \left( \frac{(x_1 + x_2 + \dots + x_m + 1)(\overline{x_1} + \dots + \overline{x_m} + 1)}{4} \right) \\ &\equiv p \sum_{x_1, \dots, x_m \in \mathbf{F}_p^\times} A \left( p, \frac{(x_1 + \dots + x_m + 1)(\overline{x_1} + \dots + \overline{x_m} + 1)}{4} \right) \pmod{p^2}. \end{aligned}$$

Proposition 3.1 shows that the moments  $S(k)$  vanish modulo  $p$ , hence writing  $T_n$  as a linear combination of moments gives  $T_n \equiv (-1)^n S(n) \pmod{p^2}$ .  $\square$

**Remark.** *The number of simultaneous solutions to the congruences  $x_1 + x_2 + \dots + x_p = a$  and  $\overline{x_1} + \overline{x_2} + \dots + \overline{x_p} = b$  is equal to  $0 \pmod{p}$  for all  $(a, b)$  except for  $a = b = 0$ , where the number of solutions is  $-1 \pmod{p}$ . Applying this fact yields that  $T_{m+p} \equiv -T_m \pmod{p^2}$ .*

#### ACKNOWLEDGEMENTS

The authors would like to thank Ken Ono, Amanda Folsom, and Riad Masri for running the UW-Madison REU in Number Theory. We would also like to thank Ken Ono in particular for his guidance in preparing this manuscript and his persistent encouragement, and also the anonymous referee for their helpful comments.

#### REFERENCES

- [1] S. Ahlgren, *Gaussian hypergeometric series and combinatorial congruences*, Symbolic computation, number theory, special functions, physics and combinatorics [Ed. F. Garvan and M. E. H. Ismail, Kluwer Acad. Publ. Dordrecht (2001), 1-12.
- [2] S. Ahlgren, K. Ono, *Modularity of a certain Calabi-Yau threefold*, Monatsh. Math. 129 (2000), 177-190
- [3] S. Ahlgren, K. Ono, *A Gaussian hypergeometric series evaluation and Apéry number congruences*, J. Reine Angew. Math 518 (2000) 187-212
- [4] R. Apéry, *Irrationalité de  $\zeta(2)$  et  $\zeta(3)$* , Astérisque 61 (1979), 11-13
- [5] R. J. Evans, *Hypergeometric  ${}_3F_2(1/4)$  evaluations over finite fields and Hecke eigenforms*, Proc. Amer. Math. Soc. to appear.
- [6] R. J. Evans, and J. Greene, *Clausen's theorem and hypergeometric functions over finite fields*, Finite Fields Appl. 15 (2009)97-109
- [7] J. Greene, *Hypergeometric functions over finite fields*, Trans. Amer. Math. Soc. 301 (1987), 77-101
- [8] K. Ireland and M. Rosen, *A classical introduction to modern number theory*, Springer-Verlag, New York, 1990.
- [9] D. H. Lehmer, Emma Lehmer, *On the cubes of Kloosterman sums*, Acta Arith. 6 (1960), 15-22
- [10] L. J. Mordell, *On Lehmer's Congruence Associated with Cubes of Kloosterman's Sums*, Journal London Math. Soc. 36 (1961), 335-339
- [11] K. Ono, *The Web of Modularity: Arithmetic of the Coefficients of Modular Forms and  $q$ -Series*, CBMS, vol. 102, Amer. Math. Soc., Providence, RI, 2004.

673 MT. PHILO RD., SHELBURNE, VT 05482

*E-mail address:* Dummit@math.wisc.edu

617 LOGAN LN., DANVILLE, CA 94526

*E-mail address:* AdamWGoldberg@gmail.com

3301 HIGHLAND AVE., DREXEL HILL, PA 19026

*E-mail address:* arp2125@columbia.edu