



Combinatorics in affine flag varieties

James Parkinson^a, Arun Ram^{b,c,*}, Christoph Schwer^d

^a *Institut für Mathematische Strukturtheorie, Technische Universität Graz, Steyrergasse 30/III, A-8010 Graz, Austria*

^b *Department of Mathematics and Statistics, University of Melbourne, Parkville, VIC 3010, Australia*

^c *Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA*

^d *Mathematisches Institut, Universität zu Köln, Weyertal 86-90, 50931 Köln, Germany*

Received 4 January 2008

Communicated by Andrew Mathas

Dedicated to Gus Lehrer on the occasion of his 60th birthday

Abstract

The Littelmann path model gives a realization of the crystals of integrable representations of symmetrizable Kac–Moody Lie algebras. Recent work of Gaussett and Littelmann [S. Gaussett, P. Littelmann, LS galleries, the path model, and MV cycles, *Duke Math. J.* 127 (1) (2005) 35–88] and others [A. Braverman, D. Gaijsgory, Crystals via the affine Grassmannian, *Duke Math. J.* 107 (3) (2001) 561–575; S. Gaussett, G. Rousseau, Kac–Moody groups, hovels and Littelmann’s paths, preprint, arXiv:math.GR/0703639, 2007] has demonstrated a connection between this model and the geometry of the loop Grassmannian. The alcove walk model is a version of the path model which is intimately connected to the combinatorics of the affine Hecke algebra. In this paper we define a refined alcove walk model which encodes the points of the affine flag variety. We show that this combinatorial indexing naturally indexes the cells in generalized Mirković–Vilonen intersections.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Loop Grassmannian; Path model; MV cycles

* Corresponding author at: Department of Mathematics and Statistics, University of Melbourne, Parkville, VIC 3010, Australia.

E-mail addresses: parkinson@weyl.math.tu-graz.ac.at (J. Parkinson), a.ram@ms.unimelb.edu.au (A. Ram), cschwer@math.uni-koeln.de (C. Schwer).

1. Introduction

A Chevalley group is a group in which row reduction works. This means that it is a group with a special set of generators (the “elementary matrices”) and relations which are generalizations of the usual row reduction operations. One way to efficiently encode these generators and relations is with a Kac–Moody Lie algebra \mathfrak{g} . From the data of the Kac–Moody Lie algebra and a choice of a commutative ring or field \mathbb{F} the group $G(\mathbb{F})$ is built by generators and relations following Chevalley–Steinberg–Tits.

Of particular interest is the case where \mathbb{F} is the field of fractions of \mathfrak{o} , the discrete valuation ring \mathfrak{o} is the ring of integers in \mathbb{F} , \mathfrak{p} is the unique maximal ideal in \mathfrak{o} and $k = \mathfrak{o}/\mathfrak{p}$ is the residue field. The favorite examples are

$$\begin{aligned} \mathbb{F} = \mathbb{C}((t)), & \quad \mathfrak{o} = \mathbb{C}[[t]], & \quad k = \mathbb{C}, \\ \mathbb{F} = \mathbb{Q}_p, & \quad \mathfrak{o} = \mathbb{Z}_p, & \quad k = \mathbb{F}_p, \\ \mathbb{F} = \mathbb{F}_q((t)), & \quad \mathfrak{o} = \mathbb{F}_q[[t]], & \quad k = \mathbb{F}_q, \end{aligned}$$

where \mathbb{Q}_p is the field of p -adic numbers, \mathbb{Z}_p is the ring of p -adic integers, and \mathbb{F}_q is the finite field with q elements. For clarity of presentation we shall work in the first case where $\mathbb{F} = \mathbb{C}((t))$. The diagram

$$\begin{array}{ccc} \mathbb{F} & & G = G(\mathbb{C}((t))) \\ \cup & & \cup \\ \mathfrak{o} \xrightarrow{\text{ev}_{t=0}} k = \mathfrak{o}/\mathfrak{p} & \text{gives} & \begin{array}{ccc} K = G(\mathbb{C}[[t]]) & \xrightarrow{\text{ev}_{t=0}} & G(\mathbb{C}) \\ \cup & & \cup \\ I = \text{ev}_{t=0}^{-1}(B(\mathbb{C})) & \xrightarrow{\text{ev}_{t=0}} & B(\mathbb{C}) \end{array} \end{array} \quad (1.1)$$

where $B(\mathbb{C})$ is the “Borel subgroup” of “upper triangular matrices” in $G(\mathbb{C})$. The loop group is $G = G(\mathbb{C}((t)))$, I is the standard Iwahori subgroup of G ,

$$\begin{aligned} G(\mathbb{C})/B(\mathbb{C}) & \text{ is the flag variety,} \\ G/I & \text{ is the affine flag variety, and } G/K \text{ is the loop Grassmanian.} \end{aligned} \quad (1.2)$$

The primary tool for the study of these varieties (ind-schemes) are the following “classical” double coset decompositions, see [St, Ch. 8] and [Mac1, §(2.6)].

Theorem 1.1. Let W be the Weyl group of $G(\mathbb{C})$, $\tilde{W} = W \times \mathfrak{h}_{\mathbb{Z}}$ the affine Weyl group, and U^- the subgroup of “unipotent lower triangular” matrices in $G(\mathbb{F})$ and $\mathfrak{h}_{\mathbb{Z}}^+$ the set of dominant elements of $\mathfrak{h}_{\mathbb{Z}}$. Then

$$\begin{array}{lll} \text{Bruhat decomposition} & G = \bigsqcup_{w \in W} BwB, & K = \bigsqcup_{w \in W} IwI, \\ \text{Iwahori decomposition} & G = \bigsqcup_{w \in \tilde{W}} IwI, & G = \bigsqcup_{v \in \tilde{W}} U^-vI, \\ \text{Cartan decomposition} & G = \bigsqcup_{\lambda^\vee \in \mathfrak{h}_{\mathbb{Z}}^+} Kt_\lambda K, & G = \bigsqcup_{\mu^\vee \in \mathfrak{h}_{\mathbb{Z}}} U^-t_\mu K, \end{array} \quad \text{Iwasawa decomposition.}$$

It should be stressed that we have, intentionally, *not* given precise definitions of the objects in Theorem 1.1. Even in the classical case, the definition of $\mathfrak{h}_{\mathbb{Z}}$ in Theorem 1.1 is sensitive to small changes in the definition of G (center, completions, etc.) and there are subtleties in making these definitions correctly in general. These issues are partly treated in [Ga1, Theorem 14.10, Lemma 6.14], [Ga2, Theorem 1.8], [GR, Remark 6.10] and [BF, Proposition 3.7].

In this paper we shall refine the Littelmann path model (in its alcove walk form, see [Ra]) by putting labels on the paths to provide a combinatorial indexing of the points in the affine flag variety. This combinatorial method of expressing the points of G/I gives detailed information about the structure of the intersections

$$U^{-}vI \cap IwI \quad \text{with } v, w \in \tilde{W}. \tag{1.3}$$

The corresponding intersections in G/K have arisen in many contexts. Most notably, the set of *Mirković–Vilonen cycles of shape λ^{\vee} and weight μ^{\vee}* is the set of irreducible components of the closure of $U^{-}t_{\mu^{\vee}}K \cap Kt_{\lambda^{\vee}}K$ in G/K ,

$$MV(\lambda^{\vee})_{\mu^{\vee}} = \text{Irr}(\overline{U^{-}t_{\mu^{\vee}}K \cap Kt_{\lambda^{\vee}}K}),$$

and

$$\text{when } k = \mathbb{F}_q, \quad \text{Card}_{G/K}(U^{-}t_{\mu^{\vee}}K \cap Kt_{\lambda^{\vee}}K) \text{ is}$$

(up to some easily understood factors) the coefficient of the monomial symmetric function $m_{\mu^{\vee}}$ in the expansion of the Macdonald spherical function $P_{\lambda^{\vee}}$.

Sections 2–6 give elementary treatments of Borchers–Kac–Moody Lie algebras, Chevalley groups, the flag variety, loop groups and affine flag varieties. With future developments in mind we have presented this material in the context of loop groups of symmetrizable Kac–Moody groups. In spite of the generality in Sections 2–6, the main results of this paper, given in Section 7, are only for loop groups of finite dimensional Chevalley groups. We do have some results in the more general case, but the restrictions of time and space have forced us to postpone the exposition of these results to a future paper.

2. Borchers–Kac–Moody Lie algebras

This section reviews definitions and sets notations for Borchers–Kac–Moody Lie algebras. Standard references are the book of Kac [Kac], the books of Wakimoto [Wak1, Wak2], the survey article of Macdonald [Mac3] and the handwritten notes of Macdonald [Mac2]. Specifically, [Kac, Ch. 1] is a reference for Section 2.1, [Kac, Chs. 3 and 5] for Section 2.2, and [Kac, Ch. 2] for Section 2.3.

2.1. Constructing a Lie algebra from a matrix

Let $A = (a_{ij})$ be an $n \times n$ matrix. Let

$$r = \text{rank}(A), \quad \ell = \text{corank}(A), \quad \text{so that } r + \ell = n. \tag{2.1}$$

By rearranging rows and columns we may assume that $(a_{ij})_{1 \leq i, j \leq r}$ is nonsingular. Define a \mathbb{C} -vector space

$$\mathfrak{h} = \mathfrak{h}' \oplus \mathfrak{d}, \quad \text{where } \mathfrak{h}' \text{ has basis } h_1, \dots, h_n, \text{ and } \mathfrak{d} \text{ has basis } d_1, \dots, d_{\ell}. \tag{2.2}$$

Define $\alpha_1, \dots, \alpha_n \in \mathfrak{h}^*$ by

$$\alpha_i(h_j) = a_{ij} \quad \text{and} \quad \alpha_i(d_j) = \delta_{i,r+j}, \tag{2.3}$$

and let

$$\bar{\mathfrak{h}}' = \mathfrak{h}'/\mathfrak{c}, \quad \text{where } \mathfrak{c} = \{h \in \mathfrak{h}' \mid \alpha_i(h) = 0 \text{ for all } 1 \leq i \leq n\}. \tag{2.4}$$

Let $c_1, \dots, c_\ell \in \mathfrak{h}'$ be a basis of \mathfrak{c} so that $h_1, \dots, h_r, c_1, \dots, c_\ell, d_1, \dots, d_\ell$ is another basis of \mathfrak{h} and define $\kappa_1, \dots, \kappa_\ell \in \mathfrak{h}^*$ by

$$\kappa_i(h_j) = 0, \quad \kappa_i(c_j) = \delta_{ij}, \quad \text{and} \quad \kappa_i(d_j) = 0. \tag{2.5}$$

Then $\alpha_1, \dots, \alpha_n, \kappa_1, \dots, \kappa_\ell$ form a basis of \mathfrak{h}^* . Let \mathfrak{a} be the Lie algebra given by generators $\mathfrak{h}, e_1, \dots, e_n, f_1, \dots, f_n$ and relations

$$[h, h'] = 0, \quad [e_i, f_j] = \delta_{ij}h_i, \quad [h, e_i] = \alpha_i(h)e_i, \quad [h, f_i] = -\alpha_i(h)f_i, \tag{2.6}$$

for $h, h' \in \mathfrak{h}$ and $1 \leq i, j \leq n$. The *Borcherds–Kac–Moody Lie algebra* of A is

$$\mathfrak{g} = \frac{\mathfrak{a}}{\mathfrak{t}}, \quad \text{where } \mathfrak{t} \text{ is the largest ideal of } \mathfrak{a} \text{ such that } \mathfrak{t} \cap \mathfrak{h} = 0. \tag{2.7}$$

The Lie algebra \mathfrak{a} is graded by

$$Q = \sum_{i=1}^n \mathbb{Z}\alpha_i, \quad \text{by setting } \deg(e_i) = \alpha_i, \deg(f_i) = -\alpha_i, \deg(h) = 0, \tag{2.8}$$

for $h \in \mathfrak{h}$. Any ideal of \mathfrak{a} is Q -graded and so \mathfrak{g} is Q -graded (see [Mac2, (1.6)] or [Mac3, p. 81]),

$$\mathfrak{g} = \mathfrak{g}_0 \oplus \left(\bigoplus_{\alpha \in R} \mathfrak{g}_\alpha \right), \quad \text{where } \mathfrak{g}_\alpha = \{x \in \mathfrak{g} \mid [h, x] = \alpha(h)x\}, \text{ and} \tag{2.9}$$

$$R = \{\alpha \mid \alpha \neq 0 \text{ and } \mathfrak{g}_\alpha \neq 0\} \text{ is the set of roots of } \mathfrak{g}.$$

The *multiplicity* of a root $\alpha \in R$ is $\dim(\mathfrak{g}_\alpha)$ and the decomposition of \mathfrak{g} in (2.9) is the decomposition of \mathfrak{g} as an \mathfrak{h} -module (under the adjoint action). If

\mathfrak{n}^+ is the subalgebra generated by e_1, \dots, e_n , and
 \mathfrak{n}^- is the subalgebra generated by f_1, \dots, f_n ,

then (see [Mac3, p. 83] or [Kac, §1.3])

$$\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+ \quad \text{and} \quad \mathfrak{h} = \mathfrak{g}_0, \quad \mathfrak{n}^+ = \bigoplus_{\alpha \in R^+} \mathfrak{g}_\alpha, \quad \mathfrak{n}^- = \bigoplus_{\alpha \in R^+} \mathfrak{g}_{-\alpha}, \tag{2.10}$$

where

$$R^+ = Q^+ \cap R \quad \text{with } Q^+ = \sum_{i=1}^n \mathbb{Z}_{\geq 0}\alpha_i. \tag{2.11}$$

Let \mathfrak{c} and \mathfrak{d} be as in (2.2) and (2.4). Then

$$\begin{aligned} \mathfrak{d} \text{ acts on } \mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}] \text{ by derivations, } \quad \mathfrak{c} = Z(\mathfrak{g}) = Z(\mathfrak{g}'), \\ \mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+ = \mathfrak{a}/\mathfrak{r} = \mathfrak{g}' \rtimes \mathfrak{d}, \\ \mathfrak{g}' = \mathfrak{n}^- \oplus \mathfrak{h}' \oplus \mathfrak{n}^+ = [\mathfrak{g}, \mathfrak{g}], \\ \tilde{\mathfrak{g}}' = \mathfrak{n}^- \oplus \tilde{\mathfrak{h}}' \oplus \mathfrak{n}^+ = \mathfrak{g}'/\mathfrak{c}, \end{aligned} \tag{2.12}$$

and \mathfrak{g}' is the universal central extension of $\tilde{\mathfrak{g}}'$ (see [Kac, Exercise 3.14]).

2.2. Cartan matrices, \mathfrak{sl}_2 subalgebras and the Weyl group

A Cartan matrix is an $n \times n$ matrix $A = (a_{ij})$ such that

$$a_{ij} \in \mathbb{Z}, \quad a_{ii} = 2, \quad a_{ij} \leq 0 \quad \text{if } i \neq j, \quad a_{ij} \neq 0 \quad \text{if and only if } a_{ji} \neq 0. \tag{2.13}$$

When A is a Cartan matrix the Lie algebra \mathfrak{g} contains many subalgebras isomorphic to \mathfrak{sl}_2 . For $1 \leq i \leq n$, the elements e_i and f_i act locally nilpotently on \mathfrak{g} (see [Mac3, p. 85] or [Mac2, (1.19)] or [Kac, Lemma 3.5]),

$$\text{span}\{e_i, f_i, h_i\} \cong \mathfrak{sl}_2, \quad \text{and} \quad \tilde{s}_i = \exp(\text{ad } e_i) \exp(-\text{ad } f_i) \exp(\text{ad } e_i) \tag{2.14}$$

is an automorphism of \mathfrak{g} (see [Kac, Lemma 3.8]). Thus \mathfrak{g} has lots of symmetry.

The simple reflections $s_i: \mathfrak{h}^* \rightarrow \mathfrak{h}^*$ and $s_i: \mathfrak{h} \rightarrow \mathfrak{h}$ are given by

$$s_i \lambda = \lambda - \lambda(h_i) \alpha_i \quad \text{and} \quad s_i h = h - \alpha_i(h) h_i, \quad \text{for } 1 \leq i \leq n, \tag{2.15}$$

$\lambda \in \mathfrak{h}^*, h \in \mathfrak{h}$, and

$$\tilde{s}_i \mathfrak{g} \alpha = \mathfrak{g}_{s_i \alpha} \quad \text{and} \quad \tilde{s}_i h = s_i h, \quad \text{for } \alpha \in R, h \in \mathfrak{h}.$$

The Weyl group W is the subgroup of $GL(\mathfrak{h}^*)$ (or $GL(\mathfrak{h})$) generated by the simple reflections. The simple reflections on \mathfrak{h} are reflections in the hyperplanes

$$\mathfrak{h}^{\alpha_i} = \{h \in \mathfrak{h} \mid \alpha_i(h) = 0\}, \quad \text{and} \quad \mathfrak{c} = \mathfrak{h}^W = \bigcap_{i=1}^n \mathfrak{h}^{\alpha_i}.$$

The representation of W on \mathfrak{h} and \mathfrak{h}^* are dual so that

$$\lambda(wh) = (w^{-1} \lambda)(h), \quad \text{for } w \in W, \lambda \in \mathfrak{h}^*, h \in \mathfrak{h}.$$

The group W is presented by generators s_1, \dots, s_n and relations

$$s_i^2 = 1 \quad \text{and} \quad \underbrace{s_i s_j s_i \cdots}_{m_{ij} \text{ factors}} = \underbrace{s_j s_i s_j \cdots}_{m_{ij} \text{ factors}} \tag{2.16}$$

for pairs $i \neq j$ such that $a_{ij}a_{ji} < 4$, where $m_{ij} = 2, 3, 4, 6$ if $a_{ij}a_{ji} = 0, 1, 2, 3$, respectively (see [Mac2, (2.12)] or [Kac, Proposition 3.13]).

The real roots of \mathfrak{g} are the elements of the set

$$R_{\text{re}} = \bigcup_{i=1}^n W\alpha_i, \quad \text{and} \quad R_{\text{im}} = R \setminus R_{\text{re}} \tag{2.17}$$

is the set of imaginary roots of \mathfrak{g} . If $\alpha = w\alpha_i$ is a real root then there is a subalgebra isomorphic to \mathfrak{sl}_2 spanned by

$$e_\alpha = \tilde{w}e_i, \quad f_\alpha = \tilde{w}f_i, \quad \text{and} \quad h_\alpha = \tilde{w}h_i, \tag{2.18}$$

and $s_\alpha = ws_iw^{-1}$ is a reflection in W acting on \mathfrak{h} and \mathfrak{h}^* by

$$s_\alpha\lambda = \lambda - \lambda(h_\alpha)\alpha \quad \text{and} \quad s_\alpha h = h - \alpha(h)h_\alpha, \quad \text{respectively.} \tag{2.19}$$

Let $\mathfrak{h}_{\mathbb{R}} = \mathbb{R}\text{-span}\{h_1, \dots, h_n, d_1, \dots, d_\ell\}$. The group W acts on $\mathfrak{h}_{\mathbb{R}}$ and the dominant chamber

$$C = \{\lambda^\vee \in \mathfrak{h}_{\mathbb{R}} \mid \langle \alpha_i, \lambda^\vee \rangle \geq 0 \text{ for all } 1 \leq i \leq n\} \tag{2.20}$$

is a fundamental domain for the action of W on the Tits cone

$$X = \bigcup_{w \in W} wC = \{h \in \mathfrak{h}_{\mathbb{R}} \mid \langle \alpha, h \rangle < 0 \text{ for a finite number of } \alpha \in R^+\}. \tag{2.21}$$

$X = \mathfrak{h}_{\mathbb{R}}$ if and only if W is finite (see [Kac, Proposition 3.12] and [Mac2, (2.14)]).

2.3. Symmetrizable matrices and invariant forms

A symmetrizable matrix is a matrix $A = (a_{ij})$ such that there exists a diagonal matrix

$$\mathcal{E} = \text{diag}(\epsilon_1, \dots, \epsilon_n), \quad \epsilon_i \in \mathbb{R}_{>0}, \quad \text{such that} \quad A\mathcal{E} \text{ is symmetric.} \tag{2.22}$$

If $\langle \cdot, \cdot \rangle: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ is a \mathfrak{g} -invariant symmetric bilinear form then

$$\langle h_i, h \rangle = \langle [e_i, f_i], h \rangle = -\langle f_i, [e_i, h] \rangle = \langle f_i, \alpha_i(h)e_i \rangle = \alpha_i(h)\langle e_i, f_i \rangle,$$

so that

$$\langle h_i, h \rangle = \alpha_i(h)\epsilon_i, \quad \text{where } \epsilon_i = \langle e_i, f_i \rangle. \tag{2.23}$$

Conversely, if A is a symmetrizable matrix then there is a nondegenerate invariant symmetric bilinear form on \mathfrak{g} determined by the formulas in (2.23) (see [Mac2, (3.12)] or [Kac, Theorem 2.2]).

If A is a Cartan matrix and $\langle \cdot, \cdot \rangle: \mathfrak{h} \times \mathfrak{h} \rightarrow \mathbb{C}$ is a W -invariant symmetric bilinear form then

$$\langle h_i, h \rangle = -\langle s_i h_i, h \rangle = -\langle h_i, s_i h \rangle = -\langle h_i, h - \alpha_i(h)h_i \rangle = -\langle h_i, h \rangle + \alpha_i(h)\langle h_i, h_i \rangle,$$

so that

$$\langle h_i, h \rangle = \alpha_i(h)\epsilon_i, \quad \text{where } \epsilon_i = \frac{1}{2}\langle h_i, h_i \rangle. \tag{2.24}$$

In particular, $\alpha_i(h_j)\epsilon_i = \langle h_i, h_j \rangle = \langle h_j, h_i \rangle = \alpha_j(h_i)\epsilon_j$ so that A is symmetrizable. Conversely, if A is a symmetrizable Cartan matrix then there is a nondegenerate W -invariant symmetric bilinear form on \mathfrak{h} determined by the formulas in (2.24) (see [Mac2, (2.26)]).

If $x_\alpha \in \mathfrak{g}_\alpha, y_\alpha \in \mathfrak{g}_{-\alpha}$ then $[x_\alpha, y_\alpha] \in [\mathfrak{g}_\alpha, \mathfrak{g}_{-\alpha}] \subseteq \mathfrak{g}_0 = \mathfrak{h}$ and $\langle h, [x_\alpha, y_\alpha] \rangle = -\langle [x_\alpha, h], y_\alpha \rangle = \alpha(h)\langle x_\alpha, y_\alpha \rangle$, so that

$$[x_\alpha, y_\alpha] = \langle x_\alpha, y_\alpha \rangle h_\alpha^\vee, \quad \text{where } \langle h, h_\alpha^\vee \rangle = \alpha(h) \text{ for all } h \in \mathfrak{h}, \tag{2.25}$$

determines $h_\alpha^\vee \in \mathfrak{h}$. If $\alpha \in R_{\text{re}}$ and $e_\alpha, f_\alpha, h_\alpha$ are as in (2.18) then

$$h_\alpha = [e_\alpha, f_\alpha] = \langle e_\alpha, f_\alpha \rangle h_\alpha^\vee \quad \text{and} \quad \langle e_\alpha, f_\alpha \rangle = \frac{1}{2}\langle h_\alpha, h_\alpha \rangle. \tag{2.26}$$

Let

$$\alpha^\vee = \langle e_\alpha, f_\alpha \rangle \alpha = \frac{1}{2}\langle h_\alpha, h_\alpha \rangle \alpha \quad \text{so that} \quad \alpha^\vee(h) = \langle h, h_\alpha \rangle. \tag{2.27}$$

Use the vector space isomorphism

$$\begin{array}{l} \mathfrak{h} \xrightarrow{\sim} \mathfrak{h}^* \\ h \mapsto \langle h, \cdot \rangle \\ h_\alpha \mapsto \alpha^\vee \\ h_\alpha^\vee \mapsto \alpha \end{array} \quad \text{to identify} \quad Q^\vee = \sum_{i=1}^n \mathbb{Z}h_i \quad \text{and} \quad Q^* = \sum_{i=1}^n \mathbb{Z}\alpha_i^\vee \tag{2.28}$$

and write

$$\langle \lambda^\vee, \mu \rangle = \mu(h_\lambda) \quad \text{if } \lambda^\vee = \lambda_1\alpha_1^\vee + \dots + \lambda_n\alpha_n^\vee \quad \text{and} \quad h_\lambda = \lambda_1h_1 + \dots + \lambda_nh_n. \tag{2.29}$$

3. Steinberg–Chevalley groups

This section gives a brief treatment of the theory of Chevalley groups. The primary reference is [St] and the extensions to the Kac–Moody case are found in [Ti].

Let A be a Cartan matrix and let R_{re} be the real roots of the corresponding Borchers–Kac–Moody Lie algebra \mathfrak{g} . Let U be the enveloping algebra of \mathfrak{g} . For each $\alpha \in R_{\text{re}}$ fix a choice of e_α in (2.18) (a choice of \tilde{w}). Use the notation

$$x_\alpha(t) = \exp(te_\alpha) = 1 + e_\alpha + \frac{1}{2!}t^2e_\alpha^2 + \frac{1}{3!}t^3e_\alpha^3 + \dots, \quad \text{in } U[[t]].$$

Then

$$x_\alpha(t)x_\alpha(u) = x_\alpha(t+u) \quad \text{in } U[[t, u]].$$

Following [Ti, 3.2], a *prenilpotent pair* is a pair of roots $\alpha, \beta \in R_{\text{re}}$ such that there exists $w, w' \in W$ with

$$w\alpha, w\beta \in R_{\text{re}}^+ \quad \text{and} \quad w'\alpha, w'\beta \in -R_{\text{re}}^+.$$

This condition guarantees that the Lie subalgebra of \mathfrak{g} generated by \mathfrak{g}_α and \mathfrak{g}_β is nilpotent. Let α, β be a prenilpotent pair and let $e_\alpha \in \mathfrak{g}_\alpha$ and $e_\beta \in \mathfrak{g}_\beta$ be as in (2.18). By [St, Lemma 15] there are unique integers $C_{\alpha\beta}^{i,j}$ such that

$$x_\alpha(t)x_\beta(u) = x_\beta(u)x_\alpha(t)x_{\alpha+\beta}(C_{\alpha,\beta}^{1,1}tu)x_{2\alpha+\beta}(C_{\alpha,\beta}^{2,1}t^2u)x_{\alpha+2\beta}(C_{\alpha,\beta}^{1,2}ut^2)\dots$$

Let \mathbb{F} be a commutative ring. The *Steinberg group*

$$\text{St is given by generators } x_\alpha(f) \text{ for } \alpha \in R_{\text{re}}, f \in \mathbb{F},$$

and relations

$$x_\alpha(f_1)x_\alpha(f_2) = x_\alpha(f_1 + f_2), \quad \text{for } \alpha \in R_{\text{re}}, \quad \text{and} \tag{3.1}$$

$$x_\alpha(f_1)x_\beta(f_2) = x_\beta(f_2)x_\alpha(f_1)x_{\alpha+\beta}(C_{\alpha,\beta}^{1,1}f_1f_2)x_{2\alpha+\beta}(C_{\alpha,\beta}^{2,1}f_1^2f_2)x_{\alpha+2\beta}(C_{\alpha,\beta}^{1,2}f_1f_2^2)\dots \tag{3.2}$$

for prenilpotent pairs α, β . In St define

$$n_\alpha(g) = x_\alpha(g)x_{-\alpha}(-g^{-1})x_\alpha(g), \quad n_\alpha = n_\alpha(1), \quad \text{and} \quad h_{\alpha^\vee}(g) = n_\alpha(g)n_\alpha^{-1}, \tag{3.3}$$

for $\alpha \in R_{\text{re}}$ and $g \in \mathbb{F}^\times$. Let $\mathfrak{h}_\mathbb{Z}$ be a \mathbb{Z} -lattice in \mathfrak{h} which is stable under the W -action and such that

$$\mathfrak{h}_\mathbb{Z} \supseteq Q^\vee, \quad \text{where } Q^\vee = \mathbb{Z}\text{-span}\{h_1, \dots, h_n\}$$

with h_1, \dots, h_n as in (2.2). With

T given by generators $h_{\lambda^\vee}(g)$ for $\lambda^\vee \in \mathfrak{h}_\mathbb{Z}, g \in \mathbb{F}^\times$, and relations

$$h_{\lambda^\vee}(g_1)h_{\lambda^\vee}(g_2) = h_{\lambda^\vee}(g_1g_2) \quad \text{and} \quad h_{\lambda^\vee}(g)h_{\mu^\vee}(g) = h_{\lambda^\vee+\mu^\vee}(g), \tag{3.4}$$

the *Tits group*

G is the group generated by St and T

with the relations coming from the third equation in (3.3) and the additional relations

$$h_{\lambda^\vee}(g)x_\alpha(f)h_{\lambda^\vee}(g)^{-1} = x_\alpha(g^{(\lambda^\vee, \alpha)}f) \quad \text{and} \quad n_i h_{\lambda^\vee}(g)n_i^{-1} = h_{s_i\lambda^\vee}(g). \tag{3.5}$$

For $\alpha, \beta \in R_{\text{re}}$ let $\epsilon_{\alpha\beta} = \pm 1$ be given by

$$\tilde{s}_\alpha(e_\beta) = \epsilon_{\alpha\beta} e_{s_\alpha\beta}, \quad \text{where } \tilde{s}_\alpha = \exp(\text{ad } e_\alpha) \exp(-\text{ad } f_\alpha) \exp(\text{ad } e_\alpha)$$

(see [CC, p. 48] and [Ti, (3.3)]). By [St, Lemma 37] (see also [Ti, §3.7(a)])

$$n_\alpha(g)x_\beta(f)n_\alpha(g)^{-1} = x_{s_\alpha\beta}(\epsilon_{\alpha\beta}g^{-(\beta, \alpha^\vee)}f), \quad h_{\lambda^\vee}(g)x_\beta(f)h_{\lambda^\vee}(g)^{-1} = x_\beta(g^{(\beta, \lambda^\vee)}f), \quad (3.6)$$

and

$$n_\alpha(g)h_{\lambda^\vee}(g')n_\alpha(g)^{-1} = h_{s_\alpha\lambda^\vee}(g'). \quad (3.7)$$

Thus G has a symmetry under the subgroup

$$N \text{ generated by } T \text{ and the } n_\alpha(g) \text{ for } \alpha \in R_{\text{re}}, g \in \mathbb{F}^\times. \quad (3.8)$$

If \mathbb{F} is big enough then N is the normalizer of T in G [St, Exercise (b), p. 36] and, by [St, Lemma 27], the homomorphism

$$\begin{aligned} N &\longrightarrow W \\ n_\alpha(g) &\longmapsto s_\alpha \end{aligned} \text{ is surjective with kernel } T. \quad (3.9)$$

Remark 3.1. (See [Ti, §3.7(b)].) If $\mathfrak{h}_{\mathbb{Z}} = \mathcal{Q}^\vee$ and the first relation of (3.5) holds in St then there is a surjective homomorphism $\psi: \text{St} \rightarrow G$. By [St, Lemma 22], the elements

$$n_\alpha h_{\lambda^\vee}(g) n_\alpha^{-1} h_{s_\alpha \lambda^\vee}(g)^{-1} \quad \text{and} \quad n_\alpha(g) n_\alpha^{-1} h_{\alpha^\vee}(g)^{-1}$$

automatically commute with each $x_\beta(f)$ so that $\ker(\psi) \subseteq Z(\text{St})$. In many cases St is the universal central extension of G (see [Ti, 3.7(c)] and [St, Theorems 10, 11, 12]).

Remark 3.2. The algebra $\mathfrak{g}' = [\mathfrak{g}, \mathfrak{g}]$ in (2.12) is generated by $e_\alpha, \alpha \in R_{\text{re}}$. A \mathfrak{g}' -module V is integrable if $e_\alpha, \alpha \in R_{\text{re}}$, act locally nilpotently so that

$$x_\alpha(c) = \exp(c e_\alpha), \quad \text{for } \alpha \in R_{\text{re}}, c \in \mathbb{C}, \quad (3.10)$$

are well defined operators on V . The Chevalley group G_V is the subgroup of $GL(V)$ generated by the operators in (3.10). To do this integrally use a Kostant \mathbb{Z} -form and choose a lattice in the module V (see [Ti, §4.3–4.4] and [St, Ch. 1]). The Kac–Moody group is the group G_{KM} generated by symbols

$$x_\alpha(c), \quad \alpha \in R_{\text{re}}, c \in \mathbb{C}, \quad \text{with relations } x_\alpha(c_1)x_\alpha(c_2) = x_\alpha(c_1 + c_2)$$

and the additional relations coming from forcing an element to be 1 if it acts by 1 on every integrable \mathfrak{g}' module. This is essentially the Chevalley group G_V for the case when V is the adjoint representation and so $G_{KM} \subseteq \text{Aut}(\mathfrak{g}')$. There are surjective homomorphisms

$$\text{St}(\mathbb{C}) \twoheadrightarrow G_{KM} \twoheadrightarrow G_V.$$

See [Kac, Exercises 3.16–3.19] and [Ti, Proposition 1].

Remark 3.3. [St, Lemma 28] In the setting of Remark 3.2 let T_V be the subgroup of G_V generated by $h_{\alpha^\vee}(g)$ for $\alpha \in R_{\text{re}}, g \in \mathbb{F}^\times$. Then

$$h_{\alpha_1^\vee}(g_1) \cdots h_{\alpha_n^\vee}(g_n) = 1 \quad \text{if and only if} \quad g_1^{(\mu, \alpha_1^\vee)} \cdots g_n^{(\mu, \alpha_n^\vee)} = 1 \quad \text{for all weights } \mu \text{ of } V,$$

$$Z(G_V) = \{h_{\alpha_1^\vee}(g_1) \cdots h_{\alpha_n^\vee}(g_n) \mid g_1^{(\beta, \alpha_1^\vee)} \cdots g_n^{(\beta, \alpha_n^\vee)} = 1 \text{ for all } \beta \in R\},$$

and if \mathbb{F} is big enough

$$T_V = \{h_{\omega_1^\vee}(g_1) \cdots h_{\omega_n^\vee}(g_n) \mid g_1, \dots, g_n \in \mathbb{F}^\times\},$$

where $\omega_1^\vee, \dots, \omega_n^\vee$ is a \mathbb{Z} -basis of the \mathbb{Z} -span of the weights of V [St, Lemma 35].

4. Labeling points of the flag variety G/B

In this section we follow [St, Ch. 8] to show that the points of the flag variety are naturally indexed by labeled walks. This is the first step in making a precise connection between the points in the flag variety and the alcove walk theory in [Ra].

Let G be a Tits group as in (3.5) over the field $\mathbb{F} = \mathbb{C}$. The *root subgroups*

$$\mathcal{X}_\alpha = \{x_\alpha(c) \mid c \in \mathbb{C}\}, \quad \text{for } \alpha \in R_{\text{re}}, \text{ satisfy } w\mathcal{X}_\beta w^{-1} = \mathcal{X}_{w\beta}, \tag{4.1}$$

for $w \in W$ and $\beta \in R_{\text{re}}$, since $h_{\alpha^\vee}(c)\mathcal{X}_\beta h_{\alpha^\vee}(c)^{-1} = \mathcal{X}_\beta$ and $n_\alpha \mathcal{X}_\beta n_\alpha^{-1} = \mathcal{X}_{s_\alpha \beta}$. As a group \mathcal{X}_α is isomorphic to \mathbb{C} (under addition).

The *flag variety* is G/B , where the subgroup

$$B \text{ is generated by } T \text{ and } x_\alpha(f) \text{ for } \alpha \in R_{\text{re}}^+, f \in \mathbb{C}. \tag{4.2}$$

Let $w \in W$. The *inversion set* of w is

$$R(w) = \{\alpha \in R_{\text{re}}^+ \mid w^{-1}\alpha \notin R_{\text{re}}^+\} \quad \text{and} \quad \ell(w) = \text{Card}(R(w)) \tag{4.3}$$

is the *length* of w . View a reduced expression $\vec{w} = s_{i_1} \cdots s_{i_\ell}$ in the generators in (2.16) as a *walk* in W starting at 1 and ending at w ,

$$1 \longrightarrow s_{i_1} \longrightarrow s_{i_1} s_{i_2} \longrightarrow \cdots \longrightarrow s_{i_1} \cdots s_{i_\ell} = w. \tag{4.4}$$

Letting $x_i(c) = x_{\alpha_i}(c)$ and $n_i = n_{\alpha_i}(1)$, the following theorem shows that

$$BwB = \{x_{i_1}(c_1)n_{i_1}^{-1}x_{i_2}(c_2)n_{i_2}^{-1} \cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1}B \mid c_1, \dots, c_\ell \in \mathbb{C}\} \tag{4.5}$$

so that the G/B -points of BwB are in bijection with labelings of the edges of the walk by complex numbers c_1, \dots, c_ℓ . The elements of $R(w)$ are

$$\beta_1 = \alpha_{i_1}, \quad \beta_2 = s_{i_1}\alpha_{i_2}, \quad \dots, \quad \beta_\ell = s_{i_1} \cdots s_{i_{\ell-1}}\alpha_{i_\ell}, \tag{4.6}$$

and the first relation in (3.6) gives

$$x_{i_1}(c_1)n_{i_1}^{-1}x_{i_2}(c_2)n_{i_2}^{-1}\cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1} = x_{\beta_1}(\pm c_1)\cdots x_{\beta_\ell}(\pm c_\ell)n_w, \tag{4.7}$$

where $n_w = n_{i_1}^{-1}\cdots n_{i_\ell}^{-1}$.

Theorem 4.1. (See [St, Theorem 15 and Lemma 43].) *Let $w \in W$ and let n_w be a representative of w in N . If*

$$R(w) = \{\beta_1, \dots, \beta_\ell\} \quad \text{then } \{x_{\beta_1}(c_1)\cdots x_{\beta_\ell}(c_\ell)n_w \mid c_1, \dots, c_\ell \in \mathbb{C}\}$$

is a set of representatives of the B -cosets in BwB .

Proof. The conceptual reason for this is that

$$\begin{aligned} BwB &= \left(\prod_{\alpha \in R_{\text{re}}^+} \mathcal{X}_\alpha\right)n_w B = n_w \left(\prod_{w^{-1}\alpha \notin R_{\text{re}}^+} \mathcal{X}_{w^{-1}\alpha}\right) \left(\prod_{w^{-1}\alpha \in R_{\text{re}}^+} \mathcal{X}_{w^{-1}\alpha}\right) B \\ &= n_w \left(\prod_{w^{-1}\alpha \notin R_{\text{re}}^+} \mathcal{X}_{w^{-1}\alpha}\right) B = \left(\prod_{\alpha \in R(w)} \mathcal{X}_\alpha\right)n_w B \\ &= \{x_{\beta_1}(c_1)\cdots x_{\beta_\ell}(c_\ell)n_w B \mid c_1, \dots, c_\ell \in \mathbb{F}\}. \end{aligned}$$

Since R_{re}^+ may be infinite there is a subtlety in the decomposition and ordering of the product of \mathcal{X}_α in the second ‘‘equality’’ and it is necessary to proceed more carefully. Choose a reduced decomposition $w = s_{i_1}\cdots s_{i_\ell}$ and let $\beta_1, \dots, \beta_\ell$ be the ordering of $R(w)$ from (4.6).

Step 1. Since $R(w) \subseteq R_{\text{re}}^+$ there is an inclusion

$$\{x_{\beta_1}(c_1)\cdots x_{\beta_\ell}(c_\ell)n_w B \mid c_1, \dots, c_\ell \in \mathbb{C}\} \subseteq BwB.$$

To prove equality proceed by induction on ℓ .

Base case: Suppose that $w = s_j$. Let $\alpha \in R_{\text{re}}^+$ and $c, d \in \mathbb{C}$. If $c = 0$ or α, α_j is a prenilpotent pair then, by relation (3.2),

$$x_\alpha(d)x_{\alpha_j}(c)n_j^{-1}B = x_{\alpha_j}(c')n_j^{-1}B, \quad \text{for some } c' \in \mathbb{C}. \tag{4.8}$$

If α, α_j is not a prenilpotent pair and $c \neq 0$ then $\alpha, -\alpha_j$ is a prenilpotent pair and, by (3.2),

$$x_\alpha(d)x_{\alpha_j}(c)n_j^{-1}B = x_\alpha(d)x_{-\alpha_j}(c^{-1})B = x_{-\alpha_j}(c^{-1})B = x_{\alpha_j}(c)n_j^{-1}B.$$

Thus $\{x_{\alpha_j}(c)n_j^{-1}B \mid c \in \mathbb{C}\}$ is B -invariant and so $Bs_jB = \{x_{\alpha_j}(c)n_j^{-1}B \mid c \in \mathbb{C}\}$.

Induction step: If $w = s_{i_1}\cdots s_{i_\ell}$ is reduced and if $\ell(ws_j) > \ell(w)$ then, by induction,

$$Bws_jB \subseteq BwB \cdot Bs_jB = \{x_{\beta_1}(c_1)\cdots x_{\beta_\ell}(c_\ell)x_{w\alpha_j}(c)n_w n_j^{-1}B \mid c_1, \dots, c_\ell, c \in \mathbb{F}\},$$

so that $Bws_jB = \{x_{\beta_1}(c_1)\cdots x_{\beta_{\ell+1}}(c_{\ell+1})n_{ws_j}B \mid c_1, \dots, c_{\ell+1} \in \mathbb{C}\}$ with $\beta_{\ell+1} = w\alpha_j$.

Step 2. Prove that $BwB = BvB$ if and only if $w = v$ by induction on $\ell(w)$.

Base case: Suppose that $\ell(w) = 0$. Then $BwB = BvB$ implies that $v \in B$ so that there is a representative n_v of v such that $n_v \in B \cap N$. Then $vR_{\text{re}}^+ \subseteq R_{\text{re}}^+$ since $n_v \mathcal{X}_\alpha n_v^{-1} = \mathcal{X}_{v\alpha} \in B$ for $\alpha \in R_{\text{re}}^+$. So $\ell(v) = 0$. Thus, by (2.16), $v = 1$.

Induction step: Assume $BwB = BvB$ and s_j is such that $\ell(ws_j) < \ell(w)$. Since $BvB \cdot Bs_jB \subseteq BvB \cup Bvs_jB$ (see [St, Lemma 25]),

$$Bws_jB \subseteq BwB \cdot Bs_jB = BvB \cdot Bs_jB \subseteq BvB \cup Bvs_jB = BwB \cup Bvs_jB.$$

Thus, by induction, $ws_j = w$ or $ws_j = vs_j$. Since $ws_j \neq w$, it follows that $w = v$.

Step 3. Let us show that if $x_{\alpha_{i_1}}(c_1)n_{i_1}^{-1} \cdots x_{\alpha_{i_\ell}}(c_\ell)n_{i_\ell}^{-1}B = x_{\alpha_{i_1}}(c'_1)n_{i_1}^{-1} \cdots x_{\alpha_{i_\ell}}(c'_\ell)n_{i_\ell}^{-1}B$, then $c_i = c'_i$ for $i = 1, 2, \dots, \ell$. The left hand side of

$$x_{\alpha_2}(c_2)n_{i_2}^{-1} \cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1}B = n_{i_1}x_{i_1}(c'_1 - c_1)n_{i_1}^{-1} \cdots x_{i_\ell}(c'_\ell)n_{i_\ell}^{-1}B$$

is in $Bs_{i_2} \cdots s_{i_\ell}B$. If $c'_1 \neq c_1$ then $n_{i_1}^{-1}x_{i_1}(c'_1 - c_1)n_{i_1} \in Bs_{i_1}B$ and the right hand side is contained in

$$n_{i_1}^{-1}x_{i_1}(c'_1 - c_1)n_{i_1}Bs_{i_2} \cdots s_{i_\ell}B \subseteq Bs_{i_1}B \cdot Bs_{i_2} \cdots s_{i_\ell}B = Bs_{i_1} \cdots s_{i_\ell}B.$$

By Step 2 this is impossible and so $c'_1 = c_1$. Then, by induction, $c'_i = c_i$ for $i = 1, 2, \dots, \ell$.

Step 4. From the definition of $R(w)$ it follows that if $\alpha, \beta \in R(w)$ and $\alpha + \beta \in R_{\text{re}}$ then $\alpha + \beta \in R(w)$ and if $\alpha, \beta \in R(w)$ then α, β form a prenilpotent pair. Thus, by [St, Lemma 17], any total order on the set $R(w)$ can be taken in the statement of the theorem. \square

Remark 4.2. Suppose that $\lambda \in \mathfrak{h}^*$ is dominant integral and $M(\lambda)$ is an (integrable) highest weight representation of G generated by a highest weight vector v_λ^+ . Then the set $BwBv_\lambda^+$ contains the vector wv_λ^+ and is contained in the sum $\bigoplus_{v \geq w\lambda} M(\lambda)_v$ of the weight spaces with weights $\geq w\lambda$. This is another way to show that if $w \neq v$ then $BwB \neq BvB$ and accomplish Step 2 in the proof of Theorem 4.1.

5. Loop Lie algebras and their extensions

This section gives a presentation of the theory of loop Lie algebras. The main lines of the theory are exactly as in the classical case (see, for example, [Mac2, §4] and [Kac, Ch. 7]) but, following recent trends (see [Ga2], [GK], [GR] and [Rou]) we treat the more general setting of the loop Lie algebra of a Kac–Moody Lie algebra.

Let \mathfrak{g}_0 be a symmetrizable Kac–Moody Lie algebra with bracket $[\cdot, \cdot]_0: \mathfrak{g}_0 \otimes \mathfrak{g}_0 \rightarrow \mathfrak{g}_0$ and invariant form $\langle \cdot, \cdot \rangle_0: \mathfrak{g}_0 \times \mathfrak{g}_0 \rightarrow \mathbb{C}$. The *loop Lie algebra* is

$$\mathfrak{g}_0[t, t^{-1}] = \mathbb{C}[t, t^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}_0 \quad \text{with bracket } [t^m x, t^n y]_0 = t^{m+n} [x, y]_0,$$

for $x, y \in \mathfrak{g}_0$. Let

$$\mathfrak{g} = \mathfrak{g}_0[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d, \quad \mathfrak{g}' = \mathfrak{g}_0[t, t^{-1}] \oplus \mathbb{C}c, \quad \bar{\mathfrak{g}}' = \mathfrak{g}_0[t, t^{-1}] = \frac{\mathfrak{g}'}{\mathbb{C}c}$$

where the bracket on \mathfrak{g} is given by

$$[t^m x, t^n y] = t^{m+n}[x, y]_0 + \delta_{m+n,0}m \langle x, y \rangle_0 c, \quad c \in \mathbb{Z}(\mathfrak{g}), \quad [d, t^m x] = mt^m x. \quad (5.1)$$

By [Kac, Exercise 7.8], \mathfrak{g}' is the universal central extension of $\tilde{\mathfrak{g}}'$. An invariant symmetric form on \mathfrak{g} is given by

$$\langle c, d \rangle = 1, \quad \langle c, t^m y \rangle = \langle d, t^m y \rangle = 0, \quad \langle c, c \rangle = \langle d, d \rangle = 0, \quad (5.2)$$

and

$$\langle t^m x, t^n y \rangle = \begin{cases} \langle x, y \rangle_0, & \text{if } m + n = 0, \\ 0, & \text{otherwise,} \end{cases} \quad (5.3)$$

for $x, y \in \mathfrak{g}_0, m, n \in \mathbb{Z}$.

Fix a Cartan subalgebra \mathfrak{h}_0 of \mathfrak{g}_0 and let

$$\mathfrak{h} = \mathfrak{h}_0 \oplus \mathbb{C}c \oplus \mathbb{C}d, \quad \mathfrak{h}' = \mathfrak{h}_0 \oplus \mathbb{C}c, \quad \tilde{\mathfrak{h}}' = \mathfrak{h}_0. \quad (5.4)$$

As in (2.2), let $h_1, \dots, h_n, d_1, \dots, d_\ell$ be a basis of \mathfrak{h}_0 and let

$$\begin{aligned} \{h_1, \dots, h_n, d_1, \dots, d_\ell, c, d\} &\text{ be a basis of } \mathfrak{h} \quad \text{and} \\ \{\omega_1, \dots, \omega_n, \delta_1, \dots, \delta_\ell, \Lambda_0, \delta\} &\text{ the dual basis in } \mathfrak{h}^* \end{aligned} \quad (5.5)$$

so that

$$\begin{aligned} \delta(\mathfrak{h}_0) = 0, \quad \delta(c) = 0, \quad \delta(d) = 1, \quad \text{and} \\ \Lambda_0(\mathfrak{h}_0) = 0, \quad \Lambda_0(c) = 1, \quad \Lambda_0(d) = 0. \end{aligned} \quad (5.6)$$

Let R be as in (2.9). As an \mathfrak{h} -module

$$\mathfrak{g} = \left(\bigoplus_{\substack{\alpha \in R \\ k \in \mathbb{Z}}} \mathfrak{g}_{\alpha+k\delta} \right) \oplus \left(\bigoplus_{k \in \mathbb{Z} \neq 0} \mathfrak{g}_{k\delta} \right) \oplus \mathfrak{h}, \quad \text{where } \mathfrak{h} = \mathfrak{h}_0 \oplus \mathbb{C}c \oplus \mathbb{C}d, \quad (5.7)$$

$$\mathfrak{g}_{\alpha+k\delta} = t^k \mathfrak{g}_\alpha, \quad \mathfrak{g}_{k\delta} = t^k \mathfrak{h}_0, \quad \text{and} \quad \tilde{R} = (R + \mathbb{Z}\delta) \cup \mathbb{Z}_{\neq 0}\delta \quad (5.8)$$

is the set of roots of \mathfrak{g} .

Let $\alpha \in R_{\text{re}}$ with $\alpha = w\alpha_i$ and fix a choice of e_α, f_α and h_α in (2.18) (choose \tilde{w}). Then

$$e_{-\alpha+k\delta} = t^k f_\alpha, \quad f_{-\alpha+k\delta} = t^{-k} e_\alpha, \quad h_{-\alpha+k\delta} = -h_\alpha + k \langle e_\alpha, f_\alpha \rangle_0 c, \quad (5.9)$$

span a subalgebra isomorphic to \mathfrak{sl}_2 . If $\mathfrak{g}_0 = \mathfrak{n}_0^- \oplus \mathfrak{h}_0 \oplus \mathfrak{n}_0^+$ is the decomposition in (2.10) and

\mathfrak{n}^+ is the subalgebra generated by \mathfrak{n}_0^+ and $e_{-\alpha+k\delta}$ for $\alpha \in R_{\text{re}}, k \in \mathbb{Z}_{>0}$, and

\mathfrak{n}^- is the subalgebra generated by \mathfrak{n}_0^- and $f_{-\alpha+k\delta}$ for $\alpha \in R_{\text{re}}, k \in \mathbb{Z}_{>0}$,

then

$$\mathfrak{g} = \mathfrak{n}^- \oplus \mathfrak{h} \oplus \mathfrak{n}^+ \quad \text{with } \mathfrak{n}^+ = \mathfrak{n}_0^+ \oplus \left(\bigoplus_{\substack{\alpha \in R \cup \{0\} \\ k \in \mathbb{Z}_{>0}}} \mathfrak{g}_{\alpha+k\delta} \right) \text{ and } \mathfrak{n}^- = \mathfrak{n}_0^- \oplus \left(\bigoplus_{\substack{\alpha \in R \cup \{0\} \\ k \in \mathbb{Z}_{<0}}} \mathfrak{g}_{\alpha+k\delta} \right).$$

The elements $e_{-\alpha+k\delta}$ and $f_{-\alpha+k\delta}$ in (5.9) act locally nilpotently on \mathfrak{g} because f_α and e_α act locally nilpotently on \mathfrak{g}_0 . Thus

$$\tilde{s}_{-\alpha+k\delta} = \exp(\text{ad } t^k f_\alpha) \exp(-\text{ad } t^{-k} e_\alpha) \exp(\text{ad } t^k f_\alpha) \tag{5.10}$$

is a well defined automorphism of \mathfrak{g} and

$$\tilde{s}_{-\alpha+k\delta} \mathfrak{g}_\beta = \mathfrak{g}_{s_{-\alpha+k\delta} \beta} \quad \text{and} \quad \tilde{s}_{-\alpha+k\delta} h = s_{-\alpha+k\delta} h, \tag{5.11}$$

for $h \in \mathfrak{h}$ and $\beta \in \tilde{R}$, where $s_{-\alpha+k\delta} : \mathfrak{h}^* \rightarrow \mathfrak{h}^*$ and $s_{-\alpha+k\delta} : \mathfrak{h} \rightarrow \mathfrak{h}$ are given by

$$s_{-\alpha+k\delta} \lambda = \lambda - \lambda(h_{-\alpha+k\delta})(-\alpha + k\delta) \quad \text{and} \quad s_{-\alpha+k\delta} h = h - (-\alpha + k\delta)(h)h_{-\alpha+k\delta}, \tag{5.12}$$

for $\lambda \in \mathfrak{h}^*$ and $h \in \mathfrak{h}$. The Weyl group of \mathfrak{g} is the subgroup of $GL(\mathfrak{h}^*)$ (or $GL(\mathfrak{h})$) generated by the reflections $s_{-\alpha+k\delta}$,

$$W_{\text{aff}} = \langle s_{-\alpha+k\delta} \mid \alpha \in R_{\text{re}}, k \in \mathbb{Z} \rangle. \tag{5.13}$$

Noting that $\mathfrak{h}^* = \mathfrak{h}_0^* \oplus \mathbb{C}\Lambda_0 \oplus \mathbb{C}\delta$ and $\mathfrak{h} = \mathfrak{h}_0 \oplus \mathbb{C}c \oplus \mathbb{C}d$, use (5.12) to compute

$$\begin{aligned} s_{-\alpha+k\delta}(\bar{\lambda}) &= \bar{\lambda} + \bar{\lambda}(h_\alpha)(-\alpha + k\delta), & s_{-\alpha+k\delta}(\bar{h}) &= \bar{h} + \alpha(\bar{h})(-h_\alpha + k\langle e_\alpha, f_\alpha \rangle_0 c), \\ s_{-\alpha+k\delta}(\ell\Lambda_0) &= \ell\Lambda_0 - k\ell\langle e_\alpha, f_\alpha \rangle_0(-\alpha + k\delta), & s_{-\alpha+k\delta}(mc) &= mc, \\ s_{-\alpha+k\delta}(m\delta) &= m\delta, & s_{-\alpha+k\delta}(\ell d) &= \ell d - k\ell(-h_\alpha + k\langle e_\alpha, f_\alpha \rangle_0 c) \end{aligned}$$

for $\bar{\lambda} \in \mathfrak{h}_0^*$, $\bar{h} \in \mathfrak{h}_0$, $m, \ell \in \mathbb{C}$. For $\alpha \in R_{\text{re}}$ and $k \in \mathbb{Z}$

$$\text{define } t_{k\alpha^\vee} \in W_{\text{aff}} \quad \text{by } s_{-\alpha+k\delta} = t_{k\alpha^\vee} s_{-\alpha}, \tag{5.14}$$

and use (2.26) and (2.27) to compute

$$\begin{aligned} t_{k\alpha^\vee}(\bar{\lambda}) &= \bar{\lambda} - \bar{\lambda}(kh_\alpha)\delta, & t_{k\alpha^\vee}(\bar{h}) &= \bar{h} - k\alpha^\vee(\bar{h})c, \\ t_{k\alpha^\vee}(\ell\Lambda_0) &= \ell\Lambda_0 + \ell k\alpha^\vee - \ell \frac{1}{2} \langle kh_\alpha, kh_\alpha \rangle_0 \delta, & t_{k\alpha^\vee}(mc) &= mc, \\ t_{k\alpha^\vee}(m\delta) &= m\delta, & t_{k\alpha^\vee}(\ell d) &= \ell d + \ell kh_\alpha - \ell \frac{1}{2} \langle kh_\alpha, kh_\alpha \rangle_0 c. \end{aligned}$$

Then $t_{k\alpha^\vee} t_{j\beta^\vee}(\bar{\lambda}) = t_{kh_\alpha}(\bar{\lambda} - \bar{\lambda}(jh_\beta)\delta) = \bar{\lambda} - \bar{\lambda}(kh_\alpha + jh_\beta)\delta$, and

$$\begin{aligned} t_{k\alpha^\vee} t_{j\beta^\vee}(\ell\Lambda_0) &= t_{k\alpha^\vee}(\ell\Lambda_0 + \ell j\beta^\vee - \ell \frac{1}{2} \langle jh_\beta, jh_\beta \rangle_0 \delta) \\ &= \ell\Lambda_0 + \ell k\alpha^\vee - \ell \frac{1}{2} \langle kh_\alpha, kh_\alpha \rangle_0 \delta + \ell j\beta^\vee - \ell j\beta^\vee(kh_\alpha)\delta - \ell \frac{1}{2} \langle jh_\beta, jh_\beta \rangle_0 \delta \\ &= \ell\Lambda_0 + \ell(k\alpha^\vee + j\beta^\vee) - \ell \frac{1}{2} \langle kh_\alpha + jh_\beta, kh_\alpha + jh_\beta \rangle_0 \delta. \end{aligned}$$

This computation shows that $t_{k\alpha^\vee}t_{j\beta^\vee} = t_{k\alpha^\vee+j\beta^\vee}$. Thus, if W_0 is the Weyl group of \mathfrak{g}_0 and $Q^* = \mathbb{Z}\text{-span}\{\alpha_1^\vee, \dots, \alpha_n^\vee\}$ then

$$W_{\text{aff}} = \{t_{\lambda^\vee}w \mid \lambda^\vee \in Q^*, w \in W_0\} \quad \text{with } t_{\lambda^\vee}t_{\mu^\vee} = t_{\lambda^\vee+\mu^\vee} \quad \text{and} \quad wt_{\lambda^\vee} = t_{w\lambda^\vee}w, \quad (5.15)$$

for $w \in W_0, \lambda^\vee, \mu^\vee \in Q^*$.

Since $\mathbb{C}\delta$ is W_{aff} -invariant, the group W_{aff} acts on $\mathfrak{h}^*/\mathbb{C}\delta$ and W_{aff} acts on the set

$$\begin{aligned} (\mathfrak{h}_0^* + \Lambda_0 + \mathbb{C}\delta)/\mathbb{C}\delta &\xrightarrow{\sim} \mathfrak{h}_0^* \\ \bar{\lambda} + \Lambda_0 + \mathbb{C}\delta &\longmapsto \bar{\lambda} \end{aligned} \quad (5.16)$$

and the W_{aff} -action on the right hand side is given by

$$s_\alpha(\bar{\lambda}) = \bar{\lambda} - \bar{\lambda}(h_\alpha)\alpha \quad \text{and} \quad t_{k\alpha^\vee}(\bar{\lambda}) = \bar{\lambda} + k\alpha^\vee, \quad \text{for } \bar{\lambda} \in \mathfrak{h}_0. \quad (5.17)$$

Here \mathfrak{h}_0^* is a set with a W_{aff} -action, the action of W_{aff} is *not linear*.

6. Loop groups and the affine flag variety G/I

This section gives a short treatment of loop groups following [St, Ch. 8] and [Mac1, §2.5 and 2.6]. This theory is currently a subject of intense research as evidenced by the work in [Ga2,GK, Rem,Rou,GR].

Let \mathfrak{g}_0 be a symmetrizable Kac–Moody Lie algebra and let $\mathfrak{h}_\mathbb{Z}$ be a \mathbb{Z} -lattice in \mathfrak{h}_0 that contains $Q^\vee = \mathbb{Z}\text{-span}\{h_1, \dots, h_n\}$.

$$\text{The loop group is the Tits group } G = G_0(\mathbb{C}((t))) \quad (6.1)$$

over the field $\mathbb{F} = \mathbb{C}((t))$. Let $K = G_0(\mathbb{C}[[t]])$ and $G_0(\mathbb{C})$ be the Tits groups of \mathfrak{g}_0 and $\mathfrak{h}_\mathbb{Z}$ over the rings $\mathbb{C}[[t]]$ and \mathbb{C} , respectively, and let $B(\mathbb{C})$ be the standard *Borel subgroup* of $G_0(\mathbb{C})$ as defined in (4.2). Let

$$U^- \text{ be the subgroup of } G \text{ generated by } x_{-\alpha}(f) \text{ for } \alpha \in R_{\text{re}}^+ \text{ and } f \in \mathbb{C}((t)), \quad (6.2)$$

and define the standard *Iwahori subgroup* I of G by

$$\begin{aligned} G &= G_0(\mathbb{C}((t))) \\ \cup I &\quad \cup \\ K &= G_0(\mathbb{C}[[t]]) \xrightarrow{\text{ev}_{t=0}} G_0(\mathbb{C}) \\ \cup I &\quad \cup \\ I &= \text{ev}_{t=0}^{-1}(B(\mathbb{C})) \xrightarrow{\text{ev}_{t=0}} B(\mathbb{C}). \end{aligned} \quad (6.3)$$

The *affine flag variety* is G/I .

For $\alpha + j\delta \in R_{\text{re}} + \mathbb{Z}\delta$ and $c \in \mathbb{C}$, define

$$x_{\alpha+j\delta}(c) = x_\alpha(ct^j) \quad \text{and} \quad t_{\lambda^\vee} = h_{\lambda^\vee}(t^{-1}), \quad (6.4)$$

and, for $c \in \mathbb{C}^\times$, define

$$n_{\alpha+j\delta}(c) = x_{\alpha+j\delta}(c)x_{-\alpha-j\delta}(-c^{-1})x_{\alpha+j\delta}(c), \tag{6.5}$$

$$n_{\alpha+j\delta} = n_{\alpha+j\delta}(1), \quad \text{and} \quad h_{(\alpha+j\delta)^\vee}(c) = n_{\alpha+j\delta}(c)n_{\alpha+j\delta}^{-1} \tag{6.6}$$

analogous to (3.3).

The group

$$\tilde{W} = \{t_{\lambda^\vee} w \mid \lambda^\vee \in \mathfrak{h}_{\mathbb{Z}}, w \in W_0\} \quad \text{with } t_{\lambda^\vee} t_{\mu^\vee} = t_{\lambda^\vee + \mu^\vee} \text{ and } w t_{\lambda^\vee} = t_{w\lambda^\vee} w, \tag{6.7}$$

acts on $\mathfrak{h}_0^* \oplus \mathbb{C}\delta$ by

$$v(\mu + k\delta) = v\mu + k\delta \quad \text{and} \quad t_{\lambda^\vee}(\mu + k\delta) = \mu + (k - \langle \lambda^\vee, \mu \rangle)\delta \tag{6.8}$$

for $v \in W_0$, $\lambda^\vee \in \mathfrak{h}_{\mathbb{Z}}$, $\mu \in \mathfrak{h}_{\mathbb{Z}}^*$, and $k \in \mathbb{Z}$. Then $n_{\alpha+j\delta}(c) = t_{-j\alpha^\vee} n_\alpha(c) = n_\alpha(ct^j)$,

$$n_\alpha x_{\beta+k\delta}(c)n_\alpha^{-1} = n_\alpha x_\beta(ct^k)n_\alpha^{-1} = x_{s_\alpha\beta}(\epsilon_{\alpha,\beta}ct^k) = x_{s_\alpha(\beta+k\delta)}(\epsilon_{\alpha,\beta}c)$$

for $\alpha \in R_{\text{re}}$, and, for $\lambda^\vee \in \mathfrak{h}_{\mathbb{Z}}$,

$$t_{\lambda^\vee} x_{\beta+k\delta}(c)t_{\lambda^\vee}^{-1} = x_{\beta+k\delta}(t^{-(\lambda^\vee, \beta)}c) = x_{t_{\lambda^\vee}(\beta+k\delta)}(c).$$

Thus the root subgroups

$$\mathcal{X}_{\alpha+j\delta} = \{x_{\alpha+j\delta}(c) \mid c \in \mathbb{C}\} \quad \text{satisfy } w\mathcal{X}_{\alpha+j\delta}w^{-1} = \mathcal{X}_{w(\alpha+j\delta)} \tag{6.9}$$

for $w \in \tilde{W}$ and $\alpha + j\delta \in R_{\text{re}} + \mathbb{Z}\delta$. These relations are a reflection of the symmetry of the group G under the group defined in (3.8):

$$\tilde{N} = N(\mathbb{C}((t))) \quad \text{generated by } n_\alpha(g), h_{\lambda^\vee}(g), \text{ for } g \in \mathbb{C}((t))^\times, \tag{6.10}$$

$\alpha \in R_{\text{re}}$, and $\lambda^\vee \in \mathfrak{h}_{\mathbb{Z}}$. The homomorphism $\tilde{N} \rightarrow W_0$ from (3.9) lifts to a surjective homomorphism (see [Mac1, p. 26 and p. 28])

$$\begin{aligned} \tilde{N} &\longrightarrow \tilde{W} \\ n_{\alpha+j\delta} &\longmapsto t_{-j\alpha^\vee} s_\alpha \quad \text{with kernel } H \text{ generated by } h_\lambda(d), d \in \mathbb{C}[[t]]^\times. \\ t_{\lambda^\vee} &\longmapsto t_{\lambda^\vee} \end{aligned}$$

Define

$$\tilde{R}_{\text{re}}^I = (R_{\text{re}}^+ + \mathbb{Z}_{\geq 0}\delta) \sqcup (-R_{\text{re}}^+ + \mathbb{Z}_{> 0}\delta) \quad \text{and} \quad \tilde{R}_{\text{re}}^U = -R_{\text{re}}^+ + \mathbb{Z}\delta \tag{6.11}$$

so that

$$\begin{aligned} \mathcal{X}_{\alpha+j\delta} \subseteq I &\quad \text{if and only if } \alpha + j\delta \in \tilde{R}_{\text{re}}^I \quad \text{and} \\ \mathcal{X}_{\alpha+j\delta} \subseteq U^- &\quad \text{if and only if } \alpha + j\delta \in \tilde{R}_{\text{re}}^U. \end{aligned} \tag{6.12}$$

Note that $\tilde{R}_{\text{re}}^I \sqcup (-\tilde{R}_{\text{re}}^I) = \tilde{R}_{\text{re}}^U \sqcup (-\tilde{R}_{\text{re}}^U) = R_{\text{re}} + \mathbb{Z}\delta$.

7. The folding algorithm and the intersections $U^{-\nu}I \cap IwI$

In this section we prove our main theorem, which gives a precise connection between the alcove walks in [Ra] and the points in the affine flag variety. The algorithm here is essentially that which is found in [BD] and, with our setup from the earlier sections, it is the ‘obvious one.’ The same method has, of course, been used in other contexts, see, for example, [C].

A special situation in the loop group theory is when \mathfrak{g}_0 is finite dimensional. In this case, the extended loop Lie algebra \mathfrak{g} defined in (5.1) is also a Kac–Moody Lie algebra. If G_0 is the Tits group of \mathfrak{g}_0 and $G = G_0(\mathbb{C}((t)))$ is the corresponding loop group then the subgroup I defined in (6.3) differs from the Borel subgroup of the Kac–Moody group G_{KM} for \mathfrak{g} only by elements of T , and the affine flag variety of G coincides with the flag variety of G_{KM} . Thus, in this case, Theorem 4.1 provides a labeling of the points of the affine flag variety.

Suppose that \mathfrak{g}_0 is a finite dimensional complex semisimple Lie algebra presented as a Kac–Moody Lie algebra with generators $e_1, \dots, e_n, f_1, \dots, f_n, h_1, \dots, h_n$ and Cartan matrix $A = (\alpha_i(h_j))_{1 \leq i, j \leq n}$. Let φ be the highest root of R (the highest weight of the adjoint representation), fix

$$e_\varphi \in \mathfrak{g}_\varphi, \quad f_\varphi \in \mathfrak{g}_{-\varphi} \quad \text{such that} \quad \langle e_\varphi, f_\varphi \rangle_0 = 1,$$

and let

$$e_0 = e_{-\varphi+\delta} = t f_\varphi, \quad f_0 = f_{-\varphi+\delta} = t^{-1} e_\varphi, \quad h_0 = [e_0, f_0] = [t x_{-\varphi}, t^{-1} x_\varphi] = -h_\varphi + c,$$

as in (5.9). The magical fact is that, in this case, $\mathfrak{g} = \mathfrak{g}_0[t, t^{-1}] \oplus \mathbb{C}c \oplus \mathbb{C}d$ is a Kac–Moody Lie algebra with generators $e_0, \dots, e_n, f_0, \dots, f_n, h_0, \dots, h_n, d$ and Cartan matrix

$$A^{(1)} = (\alpha_i(h_j))_{0 \leq i, j \leq n}, \quad \text{where } \alpha_0 = -\varphi + \delta \text{ and } h_0 = -h_\varphi + c, \tag{7.1}$$

where δ is as in (5.6) (see [Kac, Theorem 7.4]).

The *alcoves* are the open connected components of

$$\mathfrak{h}_\mathbb{R} \setminus \bigcup_{-\alpha+j\delta \in \bar{R}_{\text{re}}^+} H_{-\alpha+j\delta}, \quad \text{where } H_{-\alpha+j\delta} = \{x^\vee \in \mathfrak{h}_\mathbb{R} \mid \langle x^\vee, \alpha \rangle = j\}.$$


Under the map in (5.16) the chambers wC of the Tits cone X (see (2.20) and (2.21)) become the alcoves. Each alcove is a fundamental region for the action of W_{aff} on $\mathfrak{h}_\mathbb{R}$ given by (5.17) and W_{aff} acts simply transitively on the set of alcoves (see [Kac, Proposition 6.6]). Identify $1 \in W_{\text{aff}}$ with the *fundamental alcove*

$$A_0 = \{x^\vee \in \mathfrak{h}_\mathbb{R} \mid \langle x^\vee, \alpha_i \rangle > 0 \text{ for all } 0 \leq i \leq n\}$$

to make a bijection


$$W_{\text{aff}} \longleftrightarrow \{\text{alcoves}\}.$$

For example, when $\mathfrak{g}_0 = \mathfrak{sl}_3$,



The alcoves are the triangles and the (centers of) hexagons are the elements of Q^\vee .

Let $w \in W_{\text{aff}}$. Following the discussion in (4.4)–(4.6), a reduced expression $\vec{w} = s_{i_1} \cdots s_{i_\ell}$ is a walk starting at 1 and ending at w ,



and the points of

$$IwI = \{x_{i_1}(c_1)n_{i_1}^{-1}x_{i_2}(c_2)n_{i_2}^{-1}\cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1}I \mid c_1, \dots, c_\ell \in \mathbb{C}\} \tag{7.3}$$

are in bijection with labelings of the edges of the walk by complex numbers c_1, \dots, c_ℓ . The elements of $R(w) = \{\beta_1, \dots, \beta_\ell\}$ are the elements of \tilde{R}_{re}^I corresponding to the sequence of hyperplanes crossed by the walk.

The labeling of the hyperplanes in (7.2) is such that neighboring alcoves have

$$\begin{array}{c}
 H_{v\alpha_j} \\
 \left| \begin{array}{c} \rightarrow \\ \rightarrow \end{array} \right. \\
 v \qquad \qquad \qquad vs_j
 \end{array}
 \quad \text{with } v\alpha_j \in \tilde{R}_{re}^I \text{ if } v \text{ is closer to } 1 \text{ than } vs_j.
 \tag{7.4}$$

The *periodic orientation* (illustrated in (7.2)) is the orientation of the hyperplanes $H_{\alpha+k\delta}$ such that

- (a) 1 is on the positive side of H_α for $\alpha \in R_{re}^+$,
- (b) $H_{\alpha+k\delta}$ and H_α have parallel orientations.

This orientation is such that

$$v\alpha_j \in \tilde{R}_{re}^U \quad \text{if and only if} \quad
 \begin{array}{c}
 H_{v\alpha_j} \\
 \left| \begin{array}{c} - \\ + \end{array} \right. \\
 v \qquad \qquad \qquad vs_j
 \end{array}
 \tag{7.5}$$

Together, (7.4) and (7.5) provide a powerful combinatorics for analyzing the intersections $U^-vI \cap IwI$. We shall use the first identity in (3.3), in the form

$$x_\alpha(c)n_\alpha^{-1} = x_{-\alpha}(c^{-1})x_\alpha(-c)h_{\alpha^\vee}(c) \quad \text{(main folding law),} \tag{7.6}$$

to rewrite the points of IwI given in (7.3) as elements of U^-vI . Suppose that

$$x_{i_1}(c_1)n_{i_1}^{-1}\cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1} = x_{\gamma_1}(c'_1)\cdots x_{\gamma_\ell}(c'_\ell)n_v b, \quad \text{where } b \in I, \tag{7.7}$$

$v \in W_{\text{aff}}$ and $n_v = n_{j_1}^{-1}\cdots n_{j_k}^{-1}$ if $v = s_{i_1}\cdots s_{i_k}$ is a reduced word, and $\gamma_1, \dots, \gamma_\ell \in \tilde{R}_{re}^U$ so that $x_{\gamma_1}(c'_1)\cdots x_{\gamma_\ell}(c'_\ell) \in U^-$. Then the procedure described in (7.8)–(7.10) will compute $c'_{\ell+1} \in \mathbb{C}$, $b' \in I$, $v' \in W_{\text{aff}}$ and $\gamma_{\ell+1} \in \tilde{R}_{re}^U$ so that

$$x_{i_1}(c_1)n_{i_1}^{-1}\cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1}x_j(c)n_j^{-1} = x_{\gamma_1}(c'_1)\cdots x_{\gamma_\ell}(c'_\ell)x_{\gamma_{\ell+1}}(c_{\ell+1})n_{v'}b'.$$

Keep the notations in (7.7). Since $bx_j(c)n_j^{-1} \in Is_jI$ there are unique $\tilde{c} \in \mathbb{C}$ and $b' \in I$ such that $bx_j(c)n_j^{-1} = x_j(\tilde{c})n_j^{-1}b'$ and

$$\begin{aligned}
 x_{i_1}(c_1)n_{i_1}^{-1}\cdots x_{i_\ell}(c_\ell)n_{i_\ell}^{-1}x_j(c)n_j^{-1} &= x_{\gamma_1}(c'_1)\cdots x_{\gamma_\ell}(c'_\ell)n_v bx_j(c)n_j^{-1} \\
 &= x_{\gamma_1}(c'_1)\cdots x_{\gamma_\ell}(c'_\ell)n_v x_j(\tilde{c})n_j^{-1}b'.
 \end{aligned}$$

Case 1. If $v\alpha_j \in \tilde{R}_{re}^U$, $v \begin{array}{c} - \\ | \\ \xrightarrow{+} \\ \tilde{c} \end{array} \begin{array}{c} + \\ vs_j \end{array}$ then $x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_j(\tilde{c}) n_j^{-1} b'$ is equal to

$$x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) x_{v\alpha_j}(\pm\tilde{c}) n_{vs_j} b' \in U^- v s_j I \cap I w s_j I.$$

In this case, $\gamma_{\ell+1} = v\alpha_j$, $v' = vs_j$, and

$$v \begin{array}{c} - \\ | \\ \xrightarrow{+} \\ \tilde{c} \end{array} \begin{array}{c} + \\ vs_j \end{array} \text{ becomes } v \begin{array}{c} - \\ | \\ \xrightarrow{+} \\ \pm\tilde{c} \end{array} \begin{array}{c} + \\ vs_j \end{array}. \tag{7.8}$$

Case 2. If $v\alpha_j \notin \tilde{R}_{re}^U$ and $\tilde{c} \neq 0$, $vs_j \begin{array}{c} - \\ | \\ \xleftarrow{+} \\ \tilde{c} \end{array} \begin{array}{c} + \\ v \end{array}$, then

$$\begin{aligned} x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_{\alpha_j}(\tilde{c}) n_j^{-1} b' &= x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_{-\alpha_j}(\tilde{c}^{-1}) x_{\alpha_j}(-\tilde{c}) h_{\alpha_j^\vee}(\tilde{c}) b' \\ &= x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_{-\alpha_j}(\tilde{c}^{-1}) b'' \\ &= x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) x_{\gamma_{\ell+1}}(\pm\tilde{c}^{-1}) n_v b'' \in U^- v I \cap I w s_j I, \end{aligned}$$

where $\gamma_{\ell+1} = -v\alpha_j$ and $b'' = x_{\alpha_j}(-\tilde{c}) h_{\alpha_j^\vee}(\tilde{c}) b'$. So

$$vs_j \begin{array}{c} - \\ | \\ \xleftarrow{+} \\ \tilde{c} \end{array} \begin{array}{c} + \\ v \end{array} \text{ becomes } - \begin{array}{c} + \\ | \\ \xrightarrow{+} \\ \pm\tilde{c}^{-1} \end{array} \begin{array}{c} + \\ v \end{array}. \tag{7.9}$$

Case 3. If $v\alpha_j \notin \tilde{R}_{re}^U$ and $\tilde{c} = 0$, $vs_j \begin{array}{c} - \\ | \\ \xleftarrow{+} \\ 0 \end{array} \begin{array}{c} + \\ v \end{array}$, then

$$\begin{aligned} x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_{\alpha_j}(0) n_j^{-1} b' &= x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) n_v x_{-\alpha_j}(0) n_j^{-1} b' \\ &= x_{\gamma_1}(c'_1) \cdots x_{\gamma_\ell}(c'_\ell) x_{\gamma_{\ell+1}}(0) n_{vs_j} b' \in U^- v s_j I \cap I w s_j I, \end{aligned}$$

where $\gamma_{\ell+1} = -v\alpha_j$. So

$$vs_j \begin{array}{c} - \\ | \\ \xleftarrow{+} \\ 0 \end{array} \begin{array}{c} + \\ v \end{array} \text{ becomes } vs_j \begin{array}{c} - \\ | \\ \xleftarrow{+} \\ 0 \end{array} \begin{array}{c} + \\ v \end{array}. \tag{7.10}$$

We have proved the following theorem.

Theorem 7.1. If $w \in W_{\text{aff}}$ and $\vec{w} = s_{i_1} \cdots s_{i_\ell}$ is a minimal length walk to w define

$$\mathcal{P}(\vec{w})_v = \left\{ \begin{array}{l} \text{labeled folded paths } p \text{ of type } \vec{w} \\ \text{which end in } v \end{array} \right\} \quad \text{for } v \in W_{\text{aff}},$$

where a labeled folded path of type \vec{w} is a sequence of steps of the form

$$\begin{array}{ccc} \begin{array}{c} H_{v\alpha_j} \\ v \begin{array}{|c} \hline - \\ \hline \end{array} \begin{array}{|c} \hline + \\ \hline \end{array} \begin{array}{|c} \hline + \\ \hline \end{array} \\ \hline \begin{array}{|c} \hline c \\ \hline \end{array} \end{array} \begin{array}{c} H_{v\alpha_j} \\ - \begin{array}{|c} \hline + \\ \hline \end{array} \\ \hline \begin{array}{|c} \hline c^{-1} \\ \hline \end{array} \end{array} \begin{array}{c} H_{v\alpha_j} \\ v s_j \begin{array}{|c} \hline - \\ \hline \end{array} \begin{array}{|c} \hline + \\ \hline \end{array} \\ \hline \begin{array}{|c} \hline 0 \\ \hline \end{array} \end{array} \end{array} \quad \text{where the } k\text{th step has } j = i_k.$$

Viewing $U^{-}vI \cap IwI$ as a subset of G/I , there is a bijection

$$\mathcal{P}(\vec{w})_v \longleftrightarrow U^{-}vI \cap IwI.$$

Theorem 7.1 is a strengthening of the connection between the path model and the geometry of the affine flag variety as observed, in the case of the loop Grassmannian, in [GL] and, in terms of crystal bases, in [BG].

Remark 7.2. The paths in $\mathcal{P}(\vec{w})_v$ indicate a decomposition of $U^{-}vI \cap IwI$ into ‘‘cells,’’ where the cell associated to a nonlabeled path p is the set of points of $U^{-}vI \cap IwI$ which have the same underlying nonlabeled path. It would be very interesting to understand, combinatorially, the closure relations between these cells.

8. An example

For the group $G = SL_3(\mathbb{C}(\!(t)\!))$,

$$\begin{array}{lll} x_{\alpha_1}(c) = \begin{pmatrix} 1 & c & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, & h_{\alpha_1^\vee}(c) = \begin{pmatrix} c & 0 & 0 \\ 0 & c^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix}, & n_1 = \begin{pmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ x_{\alpha_2}(c) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & c \\ 0 & 0 & 1 \end{pmatrix}, & h_{\alpha_2^\vee}(c) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & 0 \\ 0 & 0 & c^{-1} \end{pmatrix}, & n_2 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \end{pmatrix}, \\ x_{\alpha_0}(c) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ ct & 0 & 1 \end{pmatrix}, & h_{\alpha_0^\vee}(c) = \begin{pmatrix} c^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & c \end{pmatrix}, & n_0 = \begin{pmatrix} 0 & 0 & -t^{-1} \\ 0 & 1 & 0 \\ t & 0 & 0 \end{pmatrix}. \end{array}$$

Let $w = s_2s_1s_0s_2s_0s_1s_0s_2s_0$ and $v = s_2s_1s_0s_2s_1s_2s_0$ so that

$$w = \begin{pmatrix} t^2 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & -t^{-2} & 0 \end{pmatrix} \quad \text{and} \quad v = \begin{pmatrix} 0 & -1 & 0 \\ t^2 & 0 & 0 \\ 0 & 0 & t^{-2} \end{pmatrix}.$$

We shall use Theorem 7.1 to show that the points of $IwI \cap U^{-}vI$ are

$$x_2(c_1)n_2^{-1}x_1(c_2)n_1^{-1}x_0(c_3)n_0^{-1}x_2(c_4)n_2^{-1}x_0(c_5)n_0^{-1}x_1(c_6)n_1^{-1}x_0(c_7)n_0^{-1}x_2(c_8)n_2^{-1}x_0(c_9)n_0^{-1}I,$$

with $c_1, \dots, c_9 \in \mathbb{C}$ such that

$$c_1 = 0, \quad c_2 = 0, \quad c_3 = 0, \quad c_4 = 0, \quad c_5 \neq 0, \quad c_6 = 0, \quad c_7 \neq 0, \quad c_9 = c_7^{-1}c_8. \quad (8.1)$$

Precisely,

$$x_2(0)n_2^{-1}x_1(0)n_1^{-1}x_0(0)n_0^{-1}x_2(0)n_2^{-1}x_0(c_5)n_0^{-1}x_1(0)n_1^{-1}x_0(c_7)n_0^{-1}x_2(c_8)n_2^{-1}x_0(c_7^{-1}c_8)n_0^{-1}$$

is equal to $u_9v_9b_9$, with $u_9 \in U^-$, $v_9 \in N$, $b_9 \in I$ given by


$$u_9 = \begin{pmatrix} 1 & 0 & 0 \\ c_5^{-1} - c_5^{-2}c_7^{-1}c_8t & 1 & 0 \\ c_5^{-1}c_7^{-1}t^{-2} & 0 & 1 \end{pmatrix}, \quad v_9 = \begin{pmatrix} 0 & 1 & 0 \\ -t^2 & 0 & 0 \\ 0 & 0 & t^{-2} \end{pmatrix}$$

$$b_9 = \begin{pmatrix} c_5^{-1} - c_5^{-2}c_7^{-1}c_8t & -c_5^{-2}c_7^{-1}c_8^2 & c_5^{-2}c_7^{-2}c_8^2 \\ -t^2 & c_5c_7 + c_8t & -c_5 - c_7^{-1}c_8t \\ -c_5^{-1}c_7^{-1}t^2 & -c_5^{-1}c_7^{-1}c_8t & c_7^{-1} + c_5^{-1}c_7^{-2}c_8t \end{pmatrix}, \quad (8.2)$$

so that $u_9 = x_{-\alpha_2}(d_1)x_{-\varphi}(d_2)x_{-\alpha_2-\delta}(d_3)x_{-\varphi-\delta}(d_4)x_{-\alpha_1}(d_5)x_{-\alpha_2-2\delta}(d_6)x_{-\varphi-3\delta}(d_7)x_{-\alpha_1+\delta}(d_8) \cdot x_{-\alpha_2-3\delta}(d_9)$ with

$$d_1 = d_2 = d_3 = d_4 = 0, \quad d_5 = c_5^{-1}, \quad d_6 = 0, \quad d_7 = c_5^{-1}c_7^{-1}, \quad d_8 = -c_5^{-2}c_7^{-1}c_8, \quad d_9 = 0.$$

Pictorially, the walk with labels c_1, \dots, c_9



the labeled folded path with labels d_1, \dots, d_9 .

The step by step computation is as follows:

Step 1. If $c_1 = 0$ then

$$x_2(c_1)n_2^{-1} = x_{-\alpha_2}(0)n_2^{-1} = u_1v_1b_1, \quad \text{with}$$

$$u_1 = x_{-\alpha_2}(0), \quad v_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad \text{and} \quad b_1 = 1.$$

Step 2. If $c_2 = 0$ then, since $v_1x_1(c_2)v_1^{-1} = x_\varphi(c_2)$,

$$u_1v_1b_1x_1(c_2)n_1^{-1} = u_1x_\varphi(c_2)v_1n_1^{-1}b_1 = u_1x_{-\varphi}(0)v_1n_1^{-1}b_1 = u_2v_2b_2, \quad \text{with}$$

$$u_2 = u_1x_{-\varphi}(0), \quad v_2 = v_1n_1^{-1} = \begin{pmatrix} 0 & -1 & 0 \\ 0 & 0 & -1 \\ 1 & 0 & 0 \end{pmatrix} \quad \text{and} \quad b_2 = 1.$$

Step 3. If $c_3 = 0$ then, since $v_2x_0(c_3)v_2^{-1} = x_{\alpha_2+\delta}(-c_3)$,

$$u_2v_2b_2x_0(c_3)n_0^{-1} = u_2x_{\alpha_2+\delta}(-c_3)v_2n_0^{-1}b_2 = u_2x_{-\alpha_2-\delta}(0)v_2n_0^{-1}b_2 = u_3v_3b_3, \quad \text{with}$$

$$u_3 = u_2x_{-\alpha_2-\delta}(0), \quad v_3 = v_2n_0^{-1} = \begin{pmatrix} 0 & -1 & 0 \\ t & 0 & 0 \\ 0 & 0 & t^{-1} \end{pmatrix}, \quad \text{and} \quad b_3 = 1.$$

Step 4. If $c_4 = 0$ then, since $v_3x_2(c_4)v_3^{-1} = x_{\varphi+\delta}(-c_4)$,

$$u_3v_3b_3x_2(c_4)n_2^{-1} = u_3x_{\varphi+\delta}(-c_4)v_3n_2^{-1}b_3 = u_3x_{-\varphi-\delta}(0)v_3n_2^{-1}b_3 = u_4v_4b_4, \quad \text{with}$$

$$u_4 = u_3x_{-\varphi-\delta}(0), \quad v_4 = v_3n_2^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ t & 0 & 0 \\ 0 & t^{-1} & 0 \end{pmatrix} \quad \text{and} \quad b_4 = 1.$$

Step 5. If $c_5 \neq 0$ then by the folding law and the fact that $v_4x_{-\alpha_0}(c_5^{-1})v_4^{-1} = x_{-\alpha_1}(c_5^{-1})$,

$$u_4v_4b_4x_0(c_5)n_0^{-1} = u_4v_4x_{-\alpha_0}(c_5^{-1})x_{\alpha_0}(-c_5)h_{\alpha_0^\vee}(c_5)b_4 = u_4x_{-\alpha_1}(c_5^{-1})v_4b_5 = u_5v_5b_5,$$

where

$$u_5 = u_4x_{-\alpha_1}(c_5^{-1}), \quad v_5 = v_4, \quad \text{and} \quad b_5 = x_{\alpha_0}(-c_5)h_{\alpha_0^\vee}(c_5)b_4 = \begin{pmatrix} c_5^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ -t & 0 & c_5 \end{pmatrix}.$$

Step 6. If $c_5^{-1}c_6 = 0$ (so $c_6 = 0$) then

$$u_5v_5b_5x_1(c_6)n_1^{-1} = u_5v_5x_1(c_5^{-1}c_6)n_1^{-1}b'_5 = u_5x_{-\alpha_2-2\delta}(0)v_5n_1^{-1}b'_5 = u_6v_6b_6,$$

with

$$u_6 = u_5x_{-\alpha_2-2\delta}(0), \quad v_6 = v_5n_1^{-1} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -t & 0 \\ t^{-1} & 0 & 0 \end{pmatrix} \quad \text{and} \quad b_6 = b'_5 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_5^{-1} & 0 \\ -c_6t & t & c_5 \end{pmatrix}$$

so that $b_5x_1(c_6)n_1^{-1} = x_1(c_5^{-1}c_6)n_1^{-1}b'_5$.

Step 7. If $c_5c_7 \neq 0$ then, since $v_6x_{-\alpha_0}(c)v_6^{-1} = x_{-\varphi-2\delta}(c)$,

$$\begin{aligned} u_6v_6b_6x_0(c_7)n_0^{-1} &= u_6v_6x_0(c_5c_7)n_0^{-1}b'_6 = u_6v_6x_{-\alpha_0}(c_5^{-1}c_7^{-1})x_{\alpha_0}(-c_5c_7)h_{\alpha_0^\vee}(c_5c_7)b'_6 \\ &= u_6x_{-\varphi-2\delta}(c_5^{-1}c_7^{-1})v_6b_7 = u_7v_7b_7, \end{aligned}$$

where

$$u_7 = u_6x_{-\varphi-2\delta}(c_5^{-1}c_7^{-1}), \quad v_7 = v_6, \quad \text{and}$$

$$b'_6 = \begin{pmatrix} c_5 & -1 & 0 \\ 0 & c_5^{-1} & 0 \\ 0 & 0 & 1 \end{pmatrix} \quad \text{and} \quad b_7 = x_{\alpha_0}(-c_5c_7)h_{\alpha_0^\vee}(c_5c_7)b'_6 = \begin{pmatrix} c_7^{-1} & -c_5^{-1}c_7^{-1} & 0 \\ 0 & c_5^{-1} & 0 \\ -c_5t & t & c_5c_7 \end{pmatrix},$$

so that $b_6x_0(c_7)n_0^{-1} = x_0(c_5c_7)n_0^{-1}b'_6$.

Step 8. No restrictions on $c_5^{-2}c_7^{-1}c_8$. Since $v_7x_{\alpha_2}(c)v_7^{-1} = x_{-\alpha_1+\delta}(-c)$,

$$u_7v_7b_7x_2(c_8)n_2^{-1} = u_7v_7x_2(c_5^{-2}c_7^{-1}c_8)n_2^{-1}b'_7 = u_7x_{-\alpha_1+\delta}(-c_5^{-2}c_7^{-1}c_8)v_7n_2^{-1}b'_7 = u_8v_8b_8,$$

with

$$u_8 = u_7x_{-\alpha_1+\delta}(-c_5^{-2}c_7^{-1}c_8), \quad v_8 = v_7n_2^{-1} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & t \\ t^{-1} & 0 & 0 \end{pmatrix}, \quad \text{and}$$

$$b_8 = b'_7 = \begin{pmatrix} c_7^{-1} & -c_5^{-1}c_7^{-1}c_8 & c_5^{-1}c_7^{-1} \\ -c_5t & c_5c_7 + c_8t & -t \\ -c_5^{-1}c_7^{-1}c_8t & c_5^{-2}c_7^{-1}c_8^2t & c_5^{-1} - c_5^{-2}c_7^{-1}c_8t \end{pmatrix},$$

so that $b_7x_2(c_8)n_2^{-1} = x_2(c_5^{-2}c_7^{-1}c_8)n_2^{-1}b'_7$.

Step 9. If $c_5^{-1}c_7c_9 - c_5^{-1}c_8 = 0$ (so $c_9 = c_7^{-1}c_8$) then

$$u_8v_8b_8x_0(c_9)n_0^{-1} = u_8v_8x_0(c_5^{-1}c_7c_9 - c_5^{-1}c_8)n_0^{-1}b'_8 = u_8x_{-\alpha_2-3\delta}(0)v_8n_0^{-1}b'_8 = u_9v_9b_9$$

with u_9, v_9 and b_9 as in (8.2).

Acknowledgments

The research of A. Ram and J. Parkinson was partially supported by the National Science Foundation under grant DMS-0353038 at the University of Wisconsin and the research of J. Parkinson was partly supported under the Austrian Science Fund grant FWF-P18703-N18. The research of C. Schwer was supported by a fellowship within the Postdoc-Programme of the German Academic Exchange Service (DAAD). J. Parkinson and C. Schwer thank the University of Wisconsin, Madison for hospitality. This paper was stimulated by the workshop on *Buildings and Combinatorial Representation Theory* at the American Institute of Mathematics March 26–30, 2007. We thank these institutions for support of our research.

References

- [BD] Y. Billig, M. Dyer, Decompositions of Bruhat type for the Kac–Moody groups, *Nova J. Algebra Geom.* 3 (1) (1994) 11–31.
- [BF] A. Braverman, M. Finkelberg, Pursuing the double affine Grassmannian I: Transversal slices via instantons on A_k -singularities, arXiv: 0711.2083.
- [BG] A. Braverman, D. Gaitsgory, Crystals via the affine Grassmannian, *Duke Math. J.* 107 (3) (2001) 561–575.
- [CC] R. Carter, Y. Chen, Automorphisms of affine Kac–Moody groups and related Chevalley groups over rings, *J. Algebra* 155 (1) (1993) 44–94, MR1206622 (94e:17032).
- [C] C. Curtis, A further refinement of the Bruhat decomposition, *Proc. Amer. Math. Soc.* 102 (1) (1988) 37–42.
- [Ga1] H. Garland, The arithmetic theory of loop groups, *Publ. Math. Inst. Hautes Études Sci.* 52 (1980) 181–312.
- [Ga2] H. Garland, A Cartan decomposition for p -adic loop groups, *Math. Ann.* 302 (1995) 151–175.
- [GK] D. Gaitsgory, D. Kazhdan, Representations of algebraic groups over a 2-dimensional local field, *Geom. Funct. Anal.* 14 (3) (2004) 535–574.
- [GL] S. Gaussent, P. Littelmann, LS galleries, the path model, and MV cycles, *Duke Math. J.* 127 (1) (2005) 35–88.
- [GR] S. Gaussent and G. Rousseau, Kac–Moody groups, hovels and Littelmann’s paths, preprint, arXiv: math.GR/0703639, 2007.
- [Kac] V. Kac, *Infinite Dimensional Lie Algebras*, third ed., Cambridge University Press, 1990.
- [Mac1] I.G. Macdonald, *Spherical functions on a group of p -adic type*, Publ. Ramanujan Institute No. 2, Madras, 1971.
- [Mac2] I.G. Macdonald, *Handwritten lecture notes on Kac–Moody algebras*, 1983.
- [Mac3] I.G. Macdonald, Kac–Moody algebras, in: D.J. Britten, F.W. Lemire, R.V. Moody (Eds.), *Lie Algebras and Related Topics*, in: Conference Proceedings of the Canadian Mathematical Society, vol. 5, 1986.
- [Ra] A. Ram, Alcove walks, Hecke algebras, Spherical functions, crystals and column strict tableaux, *Pure Appl. Math. Q.* 2 (4) (2006) 963–1013.
- [Rem] B. Rémy, Groupes de Kac–Moody déployés et presque déployés, *Astérisque* 277 (2002).
- [Rou] G. Rousseau, Groupes de Kac–Moody déployés sur un corps local, immeubles microaffines, *Compos. Math.* 142 (2006) 501–528.
- [St] R. Steinberg, *Lecture Notes on Chevalley Groups*, Yale University, 1967.
- [Ti] J. Tits, Uniqueness and presentation of Kac–Moody groups over fields, *J. Algebra* 105 (2) (1987) 542–573, MR0873684 (89b:17020).
- [Wak1] M. Wakimoto, *Infinite-Dimensional Lie Algebras*, Transl. Math. Monogr., vol. 195, American Mathematical Society, Providence, RI, ISBN 0-8218-2654-9, 2001, MR1793723 (2001k:17038).
- [Wak2] M. Wakimoto, *Lectures on Infinite-Dimensional Lie Algebra*, World Scientific Publishing Co., Inc., River Edge, NJ, ISBN 981-02-4128-3, 2001, MR1873994 (2003b:17033).