

Math 491 - Linear Algebra II, Fall 2015

Homework 7 - More on Diagonalization and Related Topics

Quiz on 4/5/16

Remark: Answers should be written in the following format:

A) Result.

B) If possible, the name of the method you used.

C) The computation or proof.

1. **Diagonalizable Operators on Invariant Subspaces.** Let $T : V \rightarrow V$ be a diagonalizable operator on a finite dimensional vector space V over a field \mathbb{F} . Suppose $W \subset V$ is a T -invariant subspace. Show that $T|_W$ is diagonalizable by considering the minimal polynomial $m_{T|_W}$.

Hint: Use the fact that T is diagonalizable if and only if its minimal polynomial is the product of distinct monic linear polynomials.

2. **Simultaneous Diagonalizability.** Let $S, T : V \rightarrow V$ be linear transformations. We say that S, T are simultaneously diagonalizable if there exists a direct sum decomposition $V = \bigoplus_{i=1}^k V_i$, and scalars $\lambda_i, \mu_i \in \mathbb{F}$, such that $T|_{V_i} = \lambda_i Id_{V_i}$, $S|_{V_i} = \mu_i Id_{V_i}$ for $i = 1, \dots, k$.
 - (a) Assume that S, T are diagonalizable. Show that $ST = TS$ if and only if S, T are simultaneously diagonalizable. Recall, that you showed one direction of this in a previous HW.
 - (b) Let V be a finite dimensional vector space over a field \mathbb{F} . Denote by $L(V)$ the set of linear transformations from V to itself.
 - (i) Show that $L(V)$ is an algebra over \mathbb{F} in a natural way. That is, it has a natural addition and multiplication.
 - (ii) Let $\mathcal{C} \subset L(V)$ be a subalgebra, i.e. closed under multiplication and addition, consisting of diagonalizable operators. Show that all elements of \mathcal{C} are simultaneously diagonalizable if and only if \mathcal{C} is commutative.

Here simultaneously diagonalizable means there exists a decomposition $V = \bigoplus_{i=1}^k V_i$ such that $T|_{V_i} = \lambda_{T_i} \cdot Id_{V_i}$ for any $T \in \mathcal{C}$. Recall, an algebra \mathcal{C} is commutative if for any $C_1, C_2 \in \mathcal{C}$ we have $C_1 C_2 = C_2 C_1$.

3. **Nilpotent Operators.** Let $T : V \rightarrow V$ be a linear transformation on a finite dimensional vector space V over \mathbb{F} . We say that T is nilpotent if there exists a flag of subspaces of V ,

$$\{0_V\} = V_0 \subset V_1 \subset \cdots \subset V_k = V,$$

such that $T(V_i) \subset V_{i-1}$ for all $i = 1, \dots, k$. We define the nilpotency degree of a nilpotent operator T as the smallest positive integer m such that $T^m = 0$.

- (a) Show the following operators are nilpotent by constructing a flag as above. In each case, compute the nilpotency degree of the given operator.
- (i) The derivative operator D on $\mathbb{F}_{\leq 3}[x]$, the space of polynomials over \mathbb{F} of degree at most 3.
- (ii) The operator $T_A : \mathbb{R}^5 \rightarrow \mathbb{R}^5$, defined by multiplication by

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

- (b) Show T is nilpotent if and only if $T^k = 0$ for some integer $k > 0$.
- (c) Show T is nilpotent if and only if there is a basis \mathcal{B} of V such that $[T]_{\mathcal{B}}$ is a strictly upper-triangular matrix, i.e. all entries on and below the diagonal are 0.
- (d) Let V be a vector space over \mathbb{F} . Consider a flag of the form:

$$\mathcal{F} : \{0_V\} = V_0 \subset V_1 \subset \cdots \subset V_k = V.$$

Denote by $N_{\mathcal{F}}$ the set of all linear transformations $T : V \rightarrow V$, such that $T(V_i) \subset V_{i-1}$ for $i = 1, \dots, k$. Show $N_{\mathcal{F}}$ is a subalgebra of $L(V)$, i.e. it is closed under addition and composition.

- (e) Let $\mathcal{N} \subset L(V)$ be a maximal collection of commuting nilpotent operators. Show that \mathcal{N} is a subalgebra.

4. **Generalized Eigenspaces.** Let $T : V \rightarrow V$ be a linear transformation on a finite dimensional vector space V over \mathbb{F} . Suppose that $m_T(x) = (x - \lambda_1)^{r_1} \cdots (x - \lambda_s)^{r_s}$. Denote by W_{λ_k} the generalized eigenspace of V associated to λ_k , that is, W_{λ_k} is the set of $v \in V$ such that $(T - \lambda_k \cdot Id_V)^m v = 0$ for some $m > 0$. Show that

$$W_{\lambda_k} = \ker(T - \lambda_k \cdot Id_V)^{r_k}.$$

5. **Similar Transformations.** Let $T, S : V \rightarrow V$ be two transformations of a finite dimensional vector space V over \mathbb{C} .

- (a) Suppose $\dim(V) = 2$ and that $m_T = m_S$. Is it true that T and S are similar?
- (b) Suppose $\dim(V) = 3$ and that $m_T = m_S$. Is it true that T and S are similar?
- (c) Suppose $\dim(V) = 3$ and that both $m_T = m_S$ and $p_T = p_S$. Is it true that T and S are similar?

Remark

The grader and the Lecturer will be happy to help you with the homework. Please visit office hours.

Good luck!