

## 10/26/11 – Calculus (Week 1)

**1.** A not uncommon calculus mistake is to believe that the product rule for derivatives says that  $(fg)' = f'g'$ . If  $f(x) = e^{x^2}$ , determine, with proof, whether there exists an open interval  $(a, b)$  and a nonzero function  $g$  defined on  $(a, b)$  such that this wrong product rule is true for  $x$  in  $(a, b)$ . (Putnam 1988)

**2.** Is there an infinite sequence of real numbers  $a_1, a_2, a_3, \dots$  such that

$$a_1^m + a_2^m + a_3^m + \dots = m$$

for every positive integer  $m$ ? (Putnam 2010)

**3.** Suppose that a sequence  $a_1, a_2, a_3, \dots$  satisfies  $0 < a_n \leq a_{2n} + a_{2n+1}$  for all  $n \geq 1$ . Prove that the series  $\sum_{n=1}^{\infty} a_n$  diverges. (Putnam 1994)

**4.** Functions  $f, g, h$  are differentiable on some open interval around 0 and satisfy the equations and initial conditions

$$\begin{aligned} f' &= 2f^2gh + \frac{1}{gh}, & f(0) &= 1, \\ g' &= fg^2h + \frac{4}{fh}, & g(0) &= 1, \\ h' &= 3fgh^2 + \frac{1}{fg}, & h(0) &= 1. \end{aligned}$$

Find an explicit formula for  $f(x)$ , valid in some open interval around 0. (Putnam 2009)

**5.** Find all differentiable functions  $f : \mathbb{R} \rightarrow \mathbb{R}$  such that

$$f'(x) = \frac{f(x+n) - f(x)}{n}$$

for all real numbers  $x$  and all positive integers  $n$ . (Putnam 2010)

**6.** Let  $f$  be a real function on the real line with continuous third derivative. Prove that there exists a point  $a$  such that

$$f(a) \cdot f'(a) \cdot f''(a) \cdot f'''(a) \geq 0.$$

(Putnam 1998)

7. Find all real-valued continuously differentiable functions  $f$  on the real line such that for all  $x$ ,

$$(f(x))^2 = \int_0^x [(f(t))^2 + (f'(t))^2] dt + 1990.$$

(Putnam 1990)

8. Let  $f$  be a twice-differentiable real-valued function satisfying

$$f(x) + f''(x) = -xg(x)f'(x),$$

where  $g(x) \geq 0$  for all real  $x$ . Prove that  $|f(x)|$  is bounded. (Putnam 1997)